antibodies -online.com







Images



Go to Product page

\sim	
()\/⊝	view
\circ	V I C V V

Quantity:	100 μL
Target:	MMP2
Binding Specificity:	AA 580-660
Reactivity:	Human, Mouse, Rat, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MMP2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (Paraffinembedded Sections) (IHC (p))

Product Details

Purpose:	MMP-2 Polyclonal Antibody
Immunogen:	Synthesized peptide derived from the C-terminal region of human MMP-2 at AA range: 580-660
Isotype:	lgG
Specificity:	MMP-2 Polyclonal Antibody detects endogenous levels of MMP-2 protein.
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen

Target Details

Target Details


Alternative Name:	MMP-2 (MMP2 Products)
Background:	Rabbit Anti-MMP-2 Polyclonal Antibody,MMP2, CLG4A, 72 kDa type IV collagenase, 72 kDa
	gelatinase, Gelatinase A, Matrix metalloproteinase-2, MMP-2, TBE-1,MMP2 is a member of the
	matrix metalloproteinase (MMP) gene family, that are zinc-dependent enzymes capable of
	cleaving components of the extracellular matrix and molecules involved in signal transduction.
	72 kDa type IV collagenase encoded by MMP2 is a gelatinase A, type IV collagenase, that
	contains three fibronectin type II repeats in its catalytic site that allow binding of denatured type
	IV and V collagen and elastin. Unlike most MMP family members, activation of this protein can
	occur on the cell membrane. This enzyme can be activated extracellularly by proteases, or,
	intracellulary by its S-glutathiolation with no requirement for proteolytical removal of the pro-
	domain. This protein is thought to be involved in multiple pathways including roles in the
	nervous system, endometrial menstrual breakdown, regulation of vascularization, and
	metastasis. Mutations in this gene have been associated with Winchester syndrome and
	Nodulosis-Arthropathy-Osteolysis (NAO) syndrome. Alternative splicing results in multiple
	transcript variants encoding different isoforms.,72 kDa type IV collagenase
Gene ID:	4313
UniProt:	P08253
Pathways:	Activation of Innate immune Response
Application Details	
Application Notes:	Optimal working dilutions should be determined experimentally by the investigator. Suggested
	starting dilutions are as follows: WB (1:500-1:2000), IF (1:200-1:1000), IHC-P (1:100-1:300),
	ELISA (1:20000). Not yet tested in other applications.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS containing 50 % Glycerol, 0.5 % BSA and 0.02 % Sodium Azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Stable for one year at -20°C from date of shipment. For maximum recovery of product,
	centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid


repeated freezing and thawing.

Images




Immunohistochemistry

Image 1. Immunohistochemical analysis of paraffinembedded rat heart tissue. 1, MMP-2 Polyclonal Antibody was diluted at 1:200 (4 °C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval (>98 °C, 20 min). 3, secondary antibody was diluted at 1:200 (room temperature, 30 min). Negative control was used by secondary antibody only.



Western Blotting

Image 2. Western Blot analysis of Jurkat (1), 3T3 (2), diluted at 1:1000.



Immunofluorescence

Image 3. Immunofluorescence analysis of mouse kidney tissue. 1, MMP-2 Polyclonal Antibody (red) was diluted at 1:200 (4 °C, overnight). 2, Cy3 Labeled secondary antibody was diluted at 1:300 (room temperature, 50 min). 3, Picture B: DAPI (blue) 10 min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B.

S	
	Please check the product details page for more images. Overall 6 images are available for ABIN7206177.