

Datasheet for ABIN720842 anti-UBAP1 antibody (AA 401-502) (FITC)



Overview

Overview	
Quantity:	100 μL
Target:	UBAP1
Binding Specificity:	AA 401-502
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UBAP1 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human UBAP1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Pig,Horse,Chicken
Purification:	Purified by Protein A.
Target Details	
Target:	UBAP1
Alternative Name:	UBAP1 (UBAP1 Products)

Target Details

Synonyms: MGC8710, NAG20, Nasopharyngeal carcinoma associated gene 20 protein, UAP,
UBAP, Ubiquitin associated protein 1, UBAP1_HUMAN.
Background: This gene is a member of the UBA domain family, whose members include
proteins having connections to ubiquitin and the ubiquitination pathway. The ubiquitin
associated domain is thought to be a non-covalent ubiquitin binding domain consisting of a
compact three helix bundle. This particular protein originates from a gene locus in a refined
region on chromosome 9 undergoing loss of heterozygosity in nasopharyngeal carcinoma
(NPC). Taking into account its cytogenetic location, this UBA domain family member is being
studied as a putative target for mutation in nasopharyngeal carcinomas. Multiple alternatively
spliced transcript variants encoding different isoforms have been found for this gene.
51271
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
For Research Use only
Liquid
1 μg/μL
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
50 % Glycerol.
ProClin
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
handled by trained staff only.
-20 °C
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
12 months