

Datasheet for ABIN721195
anti-MUM1 antibody (AA 401-500)



[Go to Product page](#)

1 Image

1 Publication

Overview

Quantity:	100 µL
Target:	MUM1
Binding Specificity:	AA 401-500
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MUM1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MUM1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Horse
Purification:	Purified by Protein A.

Target Details

Target:	MUM1
---------	------

Target Details

Alternative Name:	MUM1 (MUM1 Products)
Background:	<p>Synonyms: EXPAND1, FLJ14868, FLJ22283, HSPC211, Melanoma associated antigen mutated 1, Melanoma ubiquitous mutated protein, MGC131891, MGC163315, MUM 1, MUM-1, MUM1, MUM1_HUMAN, Mutated melanoma associated antigen 1, Mutated melanoma-associated antigen 1, Protein expandere, PWWP domain containing protein MUM1, PWWP domain-containing protein MUM1.</p> <p>Background: Involved in the DNA damage response pathway by contributing to the maintenance of chromatin architecture. Recruited to the vicinity of DNA breaks by TP53BP1 and plays an accessory role to facilitate damage-induced chromatin changes and promoting chromatin relaxation. Required for efficient DNA repair and cell survival following DNA damage.</p>
Gene ID:	84939
Pathways:	Chromatin Binding

Application Details

Application Notes:	WB 1:300-5000 ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C

Handling


Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Expiry Date: 12 months

Publications

Product cited in: Perez-Chacon, Adrados, Vallejo-Cremades, Lefebvre, Reed, Zapata: "Dysregulated TRAF3 and BCL2 Expression Promotes Multiple Classes of Mature Non-hodgkin B Cell Lymphoma in Mice." in: **Frontiers in immunology**, Vol. 9, pp. 3114, (2018) ([PubMed](#)).

Images



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Paraformaldehyde-fixed, paraffin embedded human kidney, Antigen retrieval by boiling in sodium citrate buffer (pH6) for 15min, Block endogenous peroxidase by 3% hydrogen peroxide for 30 minutes, Blocking buffer (normal goat serum) at 37°C for 20min, Antibody incubation with MUM1 Polyclonal Antibody at 1:400 overnight at 4°C, followed by a conjugated secondary and DAB staining.