

Datasheet for ABIN721204

anti-MUM1 antibody (AA 401-500) (HRP)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	MUM1
Binding Specificity:	AA 401-500
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MUM1 antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MUM1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Horse
Purification:	Purified by Protein A.

Target Details

Target:	MUM1
Alternative Name:	MUM1 (MUM1 Products)

Target Details

Background: Synonyms: EXPAND1, FLJ14868, FLJ22283, HSPC211, Melanoma associated antigen mutated 1, Melanoma ubiquitous mutated protein, MGC131891, MGC163315, MUM 1, MUM-1, MUM1, MUM1_HUMAN, Mutated melanoma associated antigen 1, Mutated melanoma-associated antigen 1, Protein expandere, PWWP domain containing protein MUM1, PWWP domain-containing protein MUM1.

Background: Involved in the DNA damage response pathway by contributing to the maintenance of chromatin architecture. Recruited to the vicinity of DNA breaks by TP53BP1 and plays an accessory role to facilitate damage-induced chromatin changes and promoting chromatin relaxation. Required for efficient DNA repair and cell survival following DNA damage.

Gene ID: 84939

Pathways: [Chromatin Binding](#)

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months