antibodies .- online.com





anti-LRRC15 antibody (AA 251-350) (Cy5)



Go to Product page

| \sim | | | |
|--------|-----|-----|-----|
| | N/P | r\/ | i⊢₩ |

Alternative Name:

Background:

| Quantity: | 100 μL | |
|-----------------------|--|--|
| Target: | LRRC15 | |
| Binding Specificity: | AA 251-350 | |
| Reactivity: | Mouse | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This LRRC15 antibody is conjugated to Cy5 | |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)) | |
| Product Details | | |
| Immunogen: | KLH conjugated synthetic peptide derived from human LRRC15 | |
| Isotype: | IgG | |
| Cross-Reactivity: | Mouse | |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit | |
| Purification: | Purified by Protein A. | |
| Target Details | | |
| Target: | LRRC15 | |

Synonyms: Leucine-rich repeat-containing protein 15, LIB, LRC15_HUMAN.

LRRC15 (LRRC15 Products)

Target Details

| rarget betane | | |
|---------------------|--|--|
| | Background: LRRC15 may contribute to regulation of cell-matrix adhesion interactions with respect to astrocyte recruitment around senile plaques in Alzheimer's disease brain. LRRC15 is induced by EWS-WT1(+KTS) in the tumor DSRCT and may play a role in cellular invasion. | |
| Gene ID: | 131578 | |
| Application Details | | |
| Application Notes: | IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |