

## Datasheet for ABIN7216026

# anti-Neurofibromin 1 antibody (AA 1520-1600)






| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | Neurofibromin 1 (NF1)  |
| Binding Specificity: | AA 1520-1600   |
| Reactivity:          | Human, Mouse, Rat  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This Neurofibromin 1 antibody is un-conjugated   |
| Application:         | ELISA, Western Blotting (WB), Immunofluorescence (IF)  |
| Product Details      |  |
| Purpose:             | Neurofibromin Polyclonal Antibody  |
| Immunogen:           | Synthesized peptide derived from the Internal region of human Neurofibromin at AA range:                             |
|                      | 1520-1600  |
| Isotype:             | IgG  |
| Specificity:         | Neurofibromin Polyclonal Antibody detects endogenous levels of Neurofibromin protein.                                |
| Purification:        | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen |
| Target Details       |  |
| Target:              | Neurofibromin 1 (NF1)  |
|                      |  |

## **Target Details**

| Alternative Name:   | Neurofibromin (NF1 Products)  |
|---------------------|---|
| Background:         | Rabbit Anti-Neurofibromin Polyclonal Antibody,NF1, Neurofibromin, Neurofibromatosis-related       |
|                     | protein NF-1,NF1 product appears to function as a negative regulator of the ras signal            |
|                     | transduction pathway. Mutations in this gene have been linked to neurofibromatosis type 1,        |
|                     | juvenile myelomonocytic leukemia and Watson syndrome. The mRNA for this gene is subject to        |
|                     | RNA editing (CGA>UGA->Arg1306Term) resulting in premature translation termination.                |
|                     | Alternatively spliced transcript variants encoding different isoforms have also been described    |
|                     | for this gene., Neurofibromin   |
| Molecular Weight:   | observerd band 319kDa   |
| Gene ID:            | 4763  |
| UniProt:            | P21359  |
| Pathways:           | cAMP Metabolic Process, Regulation of long-term Neuronal Synaptic Plasticity                      |
| Application Details |   |
| Application Notes:  | Optimal working dilutions should be determined experimentally by the investigator. Suggested      |
|                     | starting dilutions are as follows: WB (1:500-1:2000), IF (1:200-1:1000), ELISA (1:20000). Not yet |
|                     | tested in other applications.   |
| Comment:            | Primary Antibody  |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Concentration:      | 1 mg/mL   |
| Buffer:             | PBS containing 50 % Glycerol, 0.5 % BSA and 0.02 % Sodium Azide.                                  |
| Preservative:       | Sodium azide  |
| Precaution of Use:  | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which                     |
|                     | should be handled by trained staff only.  |
| Storage:            | -20 °C  |
| Storage Comment:    | Stable for one year at -20°C from date of shipment. For maximum recovery of product,              |
|                     | centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid        |

repeated freezing and thawing.

#### **Images**



#### **Western Blotting**

**Image 1.** Western Blot analysis of various cells using Neurofibromin Polyclonal Antibody.