antibodies .- online.com







anti-GLMN antibody (AA 151-250)





\sim			
	$ \backslash / \cap$	r\/I	$\triangle V$

Quantity:	100 μL	
Target:	GLMN	
Binding Specificity:	AA 151-250	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GLMN antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human FK506 binding protein associated protein	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Dog,Pig,Horse	
Purification:	Purified by Protein A.	
Target Details		


Target Details

larget Details			
Alternative Name:	FK506 binding protein associated protein (GLMN Products)		
Background:	Synonyms: FAP, GVM, GLML, FAP48, FAP68, FKBPAP, VMGLOM, Glomulin, FK506-binding		
	protein-associated protein, FKBP-associated protein, GLMN, GLMN_HUMAN		
	Background: Isoform 1: Regulatory component of cullin-RING-based SCF (SKP1-Cullin-F-box		
	protein) E3 ubiquitin-protein ligase complexes.). Inhibits E3 ubiquitin ligase activity by binding to		
	RBX1 (via RING domain) and inhibiting its interaction with the E2 ubiquitin-conjugating enzyme		
	CDC34. Inhibits RBX1-mediated neddylation of CUL1. Required for normal stability and normal		
	cellular levels of key components of SCF ubiquitin ligase complexes, including FBXW7, RBX1,		
	CUL1, CUL2, CUL3, CUL4A, and thereby contributes to the regulation of CCNE1 and MYC levels		
	(By similarity). Essential for normal development of the vasculature. Contributes to the		
	regulation of RPS6KB1 phosphorylation		
Gene ID:	11146		
UniProt:	Q92990		
Pathways:	Tube Formation		
Application Details			
Application Notes:	WB 1:300-5000		
	ELISA 1:500-1000		
	FCM 1:20-100		
	IHC-P 1:200-400		
	IHC-F 1:100-500		
	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		

Handling

	handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months	

Images



Western Blotting

Image 1. MCF-7 lysates probed with Anti-FK506 binding protein associated protein Polyclonal Antibody, Unconjugated at 1:5000 90min in 37°C.