antibodies - online.com







anti-EPH Receptor A6 antibody (AA 101-200)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	EPH Receptor A6 (Epha6)	
Binding Specificity:	AA 101-200	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This EPH Receptor A6 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Eph receptor A6	
Isotype:	IgG	
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	EPH Receptor A6 (Epha6)
Alternative Name:	Eph receptor A6 (Epha6 Products)

Target Details

Background:	Synonyms: Ehk 2, Ehk2, EPH homology kinase 2, Eph Receptor A6, EphA6, ephrin receptor	
	EphA6, ephrin type A receptor 6, Hek12, EPHA6_HUMAN.	
	Background: The Eph family of receptors comprises the largest known family of receptor	
	tyrosine kinases. Ligands of Eph family receptors are structurally related membrane bound	
	proteins that can be subdivided into two major subclasses, ephrin A and ephrin B. Expression o	
	Eph receptors is tissue specific and appears to be tied to developmental events. Ligands in the	
	ephrin A subclass, including the prototype family member ephrin A1 (B61), are membrane	
	associated through glycosylphosphatidyl inositol linkages, whereas ephrin B subclass consists	
	of ligands with transmembrane domains. The general role of the Eph family is in mediating	
	repulsive cell-cell interaction.	
Gene ID:	285220	
Pathways:	RTK Signaling	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	

\vdash	land	lına
	iaria	шц

Expiry Date:

12 months