
antibodies -online.com





anti-FGF11 antibody (AA 131-225)



\sim			
	N/P	r\/I	i⊢₩

Quantity:	100 μL
Target:	FGF11
Binding Specificity:	AA 131-225
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FGF11 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FGF11
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.
Target Details	

Target:	FGF11
Alternative Name:	FGF11 (FGF11 Products)

Target Details

Expiry Date:

rarget Betano		
Background:	Synonyms: FGF 11, FGF-11, FGF11, FGF11_HUMAN, FHF 3, FHF-3, FHF3, Fibroblast growth	
	factor 11, Fibroblast growth factor homologous factor 3, FLJ16061, MGC102953, MGC45269.	
	Background: Fibroblast growth factor-1 also designated basic FGF, are members of a family of	
	growth factors that stimulate proliferation of cells of mesenchymal, epithe-lial and	
	neuroectodermal origin. Additional members of the FGF family include the oncogenes FGF-3	
	(Int2) and FGF-4 (hst/Kaposi), FGF-5, FGF-6, FGF-7 (KGF), FGF-8 (AIGF), FGF-9 (GAF) and FGF-	
	10FGF-23. Members of the FGF family share 30-55 % amino acid sequence identity and similar	
	gene structure, and are capable of transforming cultured cells when overexpressed in	
	transfected cells. Cellular receptors for FGFs are members of a second multigene family	
	including four tyrosine kinases, designated Flg (FGFR-1), Bek (FGFR-L), TKF and FGFR-3.	
Gene ID:	2256	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	

12 months