

## Datasheet for ABIN723517

## anti-Adrenomedullin antibody (AA 122-140) (Biotin)



## Overview

Alternative Name:

Quantity:	100 μL
Target:	Adrenomedullin (ADM)
Binding Specificity:	AA 122-140
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Adrenomedullin antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Product Details  Immunogen:	KLH conjugated synthetic peptide derived from human Adrenomedullin
	KLH conjugated synthetic peptide derived from human Adrenomedullin  IgG
Immunogen:	
Immunogen: Isotype:	IgG
Immunogen: Isotype: Cross-Reactivity:	IgG Rat
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity:	IgG Rat Human,Mouse,Dog,Cow,Horse

Adrenomedullin (ADM Products)

## **Target Details**

Target Type:	Hormone
Background:	Synonyms: AM, PAMP, ADM, Adrenomedullin
	Background: AM and PAMP are potent hypotensive and vasodilatator agents. Numerous
	actions have been reported most related to the physiologic control of fluid and electrolyte
	homeostasis. In the kidney, am is diuretic and natriuretic, and both am and pamp inhibit
	aldosterone secretion by direct adrenal actions. In pituitary gland, both peptides at
	physiologically relevant doses inhibit basal ACTH secretion. Both peptides appear to act in brain
	and pituitary gland to facilitate the loss of plasma volume, actions which complement their
	hypotensive effects in blood vessels.
Gene ID:	133
UniProt:	P35318
Pathways:	Hormone Transport, Hormone Activity, C21-Steroid Hormone Metabolic Process, cAMP
	Metabolic Process, Myometrial Relaxation and Contraction, Regulation of G-Protein Coupled
	Receptor Protein Signaling, Tube Formation
Application Details	
Application Notes:	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months