

Datasheet for ABIN723644

anti-Tyrosine Hydroxylase antibody (AA 101-165) (HRP)



Go to Product page	
--------------------	--

Overview	
Quantity:	100 μL
Target:	Tyrosine Hydroxylase (TH)
Binding Specificity:	AA 101-165
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Tyrosine Hydroxylase antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human Tyrosine Hydroxylase
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow
Purification:	Purified by Protein A.
Target Details	
Target:	Tyrosine Hydroxylase (TH)
Alternative Name:	Tyrosine Hydroxylase (TH Products)

Target Details

Background:	Synonyms: TYH, DYT14, DYT5b, Tyrosine 3-monooxygenase, Tyrosine 3-hydroxylase, TH Background: Neuronal Marker Tyrosine hydroxylase is involved in the conversion of tyrosine to dopamine. As the rate-limiting enzyme in the synthesis of catecholamines, tyrosine hydroxylase has a key role in the physiology of adrenergic neurons. Tyrosine hydroxylase is regularly used as a marker for dopaminergic neurons, which is particularly relevant for research into Parkinson's disease.
Gene ID:	7054
UniProt:	P07101
Pathways:	Dopaminergic Neurogenesis, Response to Water Deprivation, Sensory Perception of Sound, Carbohydrate Homeostasis, Feeding Behaviour
Application Details	
Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months