

Datasheet for ABIN723755

anti-Presenilin 1 antibody (AA 10-80)[Go to Product page](#)**1** Publication

Overview

Quantity:	100 µL
Target:	Presenilin 1 (PSEN1)
Binding Specificity:	AA 10-80
Reactivity:	Mouse
Host:	Mouse
Clonality:	Polyclonal
Conjugate:	This Presenilin 1 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from rat presenilin 1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	Presenilin 1 (PSEN1)
---------	----------------------

Target Details

Alternative Name:	presenilin 1 (PSEN1 Products)
Background:	<p>Synonyms: Presenilin-1, PS-1, Protein S182, Psen1, Psnl1</p> <p>Background: Probable catalytic subunit of the gamma-secretase complex, an endoprotease complex that catalyzes the intramembrane cleavage of integral membrane proteins such as Notch receptors and APP (beta-amyloid precursor protein). Requires the other members of the gamma-secretase complex to have a protease activity. May play a role in intracellular signaling and gene expression or in linking chromatin to the nuclear membrane. Stimulates cell-cell adhesion though its association with the E-cadherin/catenin complex. Under conditions of apoptosis or calcium influx, cleaves E-cadherin promoting the disassembly of the E-cadherin/catenin complex and increasing the pool of cytoplasmic beta-catenin, thus negatively regulating Wnt signaling. May also play a role in hematopoiesis (By similarity).</p>
Gene ID:	29192
UniProt:	P97887
Pathways:	Notch Signaling , EGFR Signaling Pathway , Synaptic Vesicle Exocytosis , Dicarboxylic Acid Transport

Application Details

Application Notes:	<p>ELISA 1:500-1000</p> <p>IHC-P 1:200-400</p> <p>IHC-F 1:100-500</p> <p>IF(IHC-P) 1:50-200</p> <p>IF(IHC-F) 1:50-200</p> <p>IF(ICC) 1:50-200</p>
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage: 4 °C,-20 °C

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Expiry Date: 12 months

Publications

Product cited in: Chu, Peng, Long, Wang, Luo, Sharma, He: "Distribution and expression of Pen-2 in the central nervous system of APP/PS1 double transgenic mice." in: **Acta biochimica et biophysica Sinica**, Vol. 47, Issue 4, pp. 258-66, (2015) ([PubMed](#)).