

Datasheet for ABIN723905
anti-BTG2 antibody (AA 31-110)

5 Images

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	BTG2
Binding Specificity:	AA 31-110
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BTG2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin- embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human BTG2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Pig,Rabbit,Guinea Pig
Purification:	Purified by Protein A.

Target Details

Target:	BTG2
---------	------

Target Details

Alternative Name:	BTG2 (BTG2 Products)
Background:	<p>Synonyms: BTG2_HUMAN, Protein BTG2, BTG family member 2, NGF-inducible anti-prolervative protein PC3, pheochromacytoma cell-3, nerve growth factor-inducible anti-prolervative, B-cell translocation gene 2, BTG2, TIS21, PC3, TIS21.</p> <p>Background: BTG2 is a member of the BTG/Tob family. This family has structurally related proteins that appear to have antiproliferative properties. BTG2 is involved in the regulation of the G1/S transition of the cell cycle. It modulates transcription regulation mediated by ESR1(referenced from Entrez Gene). BTG2 expression is induced in vivo during neurogenesis, and the gene is transiently expressed in vitro in rat pheochromocytoma PC12 cells after induction of neuronal differentiation by addition of nerve growth factor (NGF), suggesting that BTG2 is functionally significant during the neuronal differentiation process (PMID: 12360398).</p>
Gene ID:	7832
UniProt:	P78543
Pathways:	MAPK Signaling

Application Details

Application Notes:	WB 1:300-5000 ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.


Handling

Storage: 4 °C, -20 °C

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Expiry Date: 12 months

Images




Immunofluorescence

Image 1.



Immunohistochemistry

Image 2. Formalin-fixed and paraffin embedded mouse heart tissue (ABIN723905) at 1: 200 followed by incubation with conjugated secondary antibody and DAB staining.



Immunohistochemistry

Image 3. Formalin-fixed and paraffin embedded mouse heart tissue labeled with (ABIN723905) Rabbit Anti-BTG2 TIS21 Polyclonal Antibody at 1: 200 followed by incubation with conjugated secondary antibody and DAB staining.

Images

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN723905.