

Datasheet for ABIN724115
anti-AGT antibody (AA 51-110)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	AGT
Binding Specificity:	AA 51-110
Reactivity:	Rat
Host:	Mouse
Clonality:	Polyclonal
Conjugate:	This AGT antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from rat AGT
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human, Mouse
Purification:	Purified by Protein A.

Target Details

Target:	AGT
---------	-----

Target Details

Alternative Name:	AGT (AGT Products)
Background:	<p>Synonyms: Ang, PAT, ANRT, AngII, Angiotensinogen, Serpin A8, Agt, Serpina8</p> <p>Background: Essential component of the renin-angiotensin system (RAS), a potent regulator of blood pressure, body fluid and electrolyte homeostasis. Angiotensin-2: acts directly on vascular smooth muscle as a potent vasoconstrictor, affects cardiac contractility and heart rate through its action on the sympathetic nervous system, and alters renal sodium and water absorption through its ability to stimulate the zona glomerulosa cells of the adrenal cortex to synthesize and secrete aldosterone. Angiotensin-3: stimulates aldosterone release. Angiotensin 1-7: is a ligand for the G-protein coupled receptor MAS1 (By similarity). Has vasodilator and antidiuretic effects (By similarity). Has an antithrombotic effect that involves MAS1-mediated release of nitric oxide from platelets.</p>
Gene ID:	24179
UniProt:	P01015
Pathways:	JAK-STAT Signaling , ACE Inhibitor Pathway , EGFR Signaling Pathway , Peptide Hormone Metabolism , Regulation of Systemic Arterial Blood Pressure by Hormones , Regulation of Lipid Metabolism by PPARalpha , Protein targeting to Nucleus , Feeding Behaviour , Monocarboxylic Acid Catabolic Process , Dicarboxylic Acid Transport , Positive Regulation of Response to DNA Damage Stimulus , Regulation of long-term Neuronal Synaptic Plasticity

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Formalin-fixed and paraffin embedded rat liver labeled with Anti AGT Polyclonal Antibody,Unconjugated (ABIN724115) at 1:200 followed by conjugation to the secondary antibody and DAB staining.