

# Datasheet for ABIN725127 anti-APP antibody (AA 381-480) (FITC)



#### Overview

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | APP  |
| Binding Specificity: | AA 381-480   |
| Reactivity:          | Human, Mouse, Rat  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This APP antibody is conjugated to FITC  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human of human APP |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Human, Mouse, Rat  |
| Predicted Reactivity: | Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit                           |
| Purification:         | Purified by Protein A.   |

## Target Details

| Target:           | APP                                      |
|-------------------|--|
| Alternative Name: | Amyloid Precursor Protein (APP Products) |

#### Target Details

| Background:         | Synonyms: AAA, AD1, PN2, ABPP, APPI, CVAP, ABETA, PN-II, CTFgamma, Amyloid beta A4              |
|---------------------|---|
|                     | protein, APP, Alzheimer disease amyloid protein, Cerebral vascular amyloid peptide, PreA4,      |
|                     | Protease nexin-II, A4   |
|                     | Background: This gene encodes a cell surface receptor and transmembrane precursor protein       |
|                     | that is cleaved by secretases to form a number of peptides. Some of these peptides are          |
|                     | secreted and can bind to the acetyltransferase complex APBB1/TIP60 to promote                   |
|                     | transcriptional activation, while others form the protein basis of the amyloid plaques found in |
|                     | the brains of patients with Alzheimer disease. Mutations in this gene have been implicated in   |
|                     | autosomal dominant Alzheimer disease and cerebroarterial amyloidosis (cerebral amyloid          |
|                     | angiopathy). Multiple transcript variants encoding several different isoforms have been found   |
|                     | for this gene. [provided by RefSeq, Jul 2008].  |
| Gene ID:            | 351   |
| UniProt:            | P05067  |
| Pathways:           | Caspase Cascade in Apoptosis, EGFR Signaling Pathway, Transition Metal Ion Homeostasis,         |
|                     | Skeletal Muscle Fiber Development, Toll-Like Receptors Cascades, Feeding Behaviour              |
| Application Details |   |
| Application Notes:  | IF(IHC-P) 1:50-200  |
|                     | IF(IHC-F) 1:50-200  |
|                     | IF(ICC) 1:50-200  |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Concentration:      | 1 μg/μL   |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and     |
|                     | 50 % Glycerol.  |
| Preservative:       | ProClin   |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be             |
|                     | handled by trained staff only.  |
| Storage:            | -20 °C  |
|                     |   |

## Handling

| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
|------------------|---|
| Expiry Date:     | 12 months   |