antibodies -online.com





anti-Reticulon 4 antibody (AA 301-400) (Biotin)



()	11/	IN	/ie	A .
	/ // 	۱ ات	/ (−	' \/\/

Quantity:	100 μL	
Target:	Reticulon 4 (RTN4)	
Binding Specificity:	AA 301-400	
Reactivity:	Human, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Reticulon 4 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from mouse RTN4 protein	
Isotype:	IgG	
Cross-Reactivity:	Human, Rat	
Predicted Reactivity:	Mouse	
Purification:	Purified by Protein A.	

Target Details

Target:	Reticulon 4 (RTN4)
Alternative Name:	Nogo-A (RTN4 Products)

Target Details

_			
Background:	Synonyms: ASY, NgA, NOGO, NSP-CL, Nogo-A, Nogo-B, Nogo-C, AA407876, AA409940,		
	AA960376, mKIAA0886, mKIAA4153, 1110020G17Rik, C130026I10Rik, Reticulon-4, Neurite		
	outgrowth inhibitor, Nogo protein, Rtn4, Kiaa0886		
	Background: Developmental neurite growth regulatory factor with a role as a negative regulator		
	of axon-axon adhesion and growth, and as a facilitator of neurite branching. Regulates neurite		
	fasciculation, branching and extension in the developing nervous system. Involved in down-		
	regulation of growth, stabilization of wiring and restriction of plasticity in the adult CNS.		
	Regulates the radial migration of cortical neurons via an RTN4R-LINGO1 containing receptor		
	complex. May inhibit BACE1 activity and amyloid precursor protein processing.		
Gene ID:	68585		
UniProt:	Q99P72		
Pathways:	Neurotrophin Signaling Pathway, Regulation of Cell Size, SARS-CoV-2 Protein Interactome		
Application Details			
Application Notes:	WB 1:300-5000		
	IHC-P 1:200-400		
	IHC-F 1:100-500		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C for 12 months.		
Expiry Date:	12 months		