# antibodies .- online.com





## anti-STOML2 antibody





#### Go to Product page

$\sim$							
	1//	$\Box$	$r \setminus$	/ [	$\bigcirc$	1	٨,

Quantity:	200 μL	
Target:	STOML2	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This STOML2 antibody is un-conjugated	
Application:	Immunohistochemistry (IHC)	

## Product Details

Immunogen:	Recombinant fusion protein of human STOML2 (NP_038470.1).	
Isotype:	IgG	
Characteristics:	Polyclonal Antibody	
Purification:	Affinity purification	

## **Target Details**

Target:	STOML2	
Alternative Name:	STOML2 (STOML2 Products)	
Background:	Human stomatin (band 7.2b) is a 31- kDa erythrocyte membrane protein of unknown function	
	but implicated in the control of ion channel permeability, mechanoreception, and lipid domain	
	organization. Stomatin (EPB72)-like 2 (STOML2,synonyms: SLP-2,HSPC108) is a 38.5- kDa	
	protein that is overall approximately 20 % similar to human stomatin. STOML2 is also present in	

#### **Target Details**

mature human erythrocytes ,but lacks a characteristic NH(2)-terminal hydrophobic domain found in other stomatin homologues. STOML2 may link stomatin or other integral membrane proteins to the peripheral cytoskeleton and thereby play a role in regulating ion channel conductances or the organization of sphingolipid and cholesterol-rich lipid rafts.

Gene ID: 30968

UniProt: Q9UJZ1

Pathways: SARS-CoV-2 Protein Interactome, The Global Phosphorylation Landscape of SARS-CoV-2

Infection


#### **Application Details**

Application Notes: IHC 1:50-1:200

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS with 0.02 % sodium azide, 50 % glycerol, pH 7.3
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Avoid freeze / thaw cycles.



#### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** Immunohistochemistry of paraffin-embedded Human breast cancer using STOML2 Polyclonal Antibody at dilution of 1:100 (40x lens).