antibodies -online.com





anti-CDC42 antibody





Go to Product page

)\/(

Quantity:	200 μL
Target:	CDC42
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CDC42 antibody is un-conjugated
Application:	Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant fusion protein of human CDC42 (NP_001782.1).
Isotype:	lgG
Characteristics:	Polyclonal Antibody
Purification:	Affinity purification

Target Details

Target:	CDC42
Alternative Name:	CDC42 (CDC42 Products)
Background:	The protein encoded by this gene is a small GTPase of the Rho-subfamily, which regulates signaling pathways that control diverse cellular functions including cell morphology, migration, endocytosis and cell cycle progression. This protein is highly similar to Saccharomyces cerevisiae Cdc 42, and is able to complement the yeast cdc42-1 mutant. The product of

Target Details

oncogene Dbl was reported to specifically catalyze the dissociation of GDP from this protein. This protein could regulate actin polymerization through its direct binding to Neural Wiskott-Aldrich syndrome protein (N-WASP), which subsequently activates Arp2/3 complex. Alternative splicing of this gene results in multiple transcript variants. Pseudogenes of this gene have been identified on chromosomes 3, 4, 5, 7, 8 and 20.

Gene ID: 998

UniProt: P60953

Pathways: MAPK Signaling, Microtubule Dynamics, RTK Signaling, WNT Signaling, TCR Signaling, EGFR

Signaling Pathway, Regulation of Actin Filament Polymerization, Regulation of Muscle Cell

Differentiation, Cell-Cell Junction Organization, Maintenance of Protein Location, Skeletal

Muscle Fiber Development, Signaling Events mediated by VEGFR1 and VEGFR2, EGFR

Downregulation, VEGF Signaling


Application Details

Application Notes: IHC 1:50-1:200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS with 0.02 % sodium azide, 50 % glycerol, pH 7.3
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Avoid freeze / thaw cycles.



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Immunohistochemistry of paraffin-embedded Rat testis using CDC42 Polyclonal Antibody at dilution of 1:100 (40x lens).