
# antibodies - online.com







# anti-GCSH antibody



**Images** 



### Overview

Quantity:	200 μL
Target:	GCSH
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunofluorescence (IF)

## **Product Details**

Immunogen:	Recombinant fusion protein of human GCSH (NP_004474.2).
Isotype:	IgG
Characteristics:	Polyclonal Antibody
Purification:	Affinity purification

## **Target Details**

Target:	GCSH
Alternative Name:	GCSH (GCSH Products)
Background:	Degradation of glycine is brought about by the glycine cleavage system, which is composed of
	four mitochondrial protein components: P protein (a pyridoxal phosphate-dependent glycine
	decarboxylase), H protein (a lipoic acid-containing protein), T protein (a tetrahydrofolate-
	requiring enzyme), and L protein (a lipoamide dehydrogenase). The protein encoded by this
	gene is the H protein, which transfers the methylamine group of glycine from the P protein to

## **Target Details**


	the T protein. Defects in this gene are a cause of nonketotic hyperglycinemia (NKH). Two
	transcript variants, one protein-coding and the other probably not protein-coding, have been
	found for this gene. Also, several transcribed and non-transcribed pseudogenes of this gene
	exist throughout the genome.
Molecular Weight:	Observed_MW: 19 kDa
	Calculated_MW: 18 kDa
Gene ID:	2653
UniProt:	P23434

## **Application Details**

Application Notes:	WB 1:500-1:2000 IF 1:50-1:200
Restrictions:	For Research Use only


# Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS with 0.02 % sodium azide, 50 % glycerol, pH 7.3
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Avoid freeze / thaw cycles.




### **Immunofluorescence**

**Image 1.** Immunofluorescence analysis of U-2 OS cells using GCSH Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



### **Western Blotting**

**Image 2.** Western blot analysis of extracts of various cell lines using GCSH Polyclonal Antibody at dilution of 1:3000.



## Immunofluorescence

**Image 3.** Confocal immunofluorescence analysis of U2OS cells using GCSH Polyclonal Antibody at dilution of 1:100 (60x lens). Blue: DAPI for nuclear staining.

Please check the product details page for more images. Overall 5 images are available for ABIN7257311.