antibodies -online.com





anti-OPRL1 antibody (AA 1-47) (Cy5)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL
Target:	OPRL1
Binding Specificity:	AA 1-47
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OPRL1 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from mouse Nociceptin receptor	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Predicted Reactivity:	Dog,Cow,Pig,Horse,Guinea Pig	
Purification:	Purified by Protein A.	

Target Details

Target:	OPRL1
Alternative Name:	Nociceptin receptor (OPRL1 Products)

Target Details

Background:	Synonyms: KOR3, ORGC, ORL1, Oprl, XOR1, morc, KOR-3, LC132, MOR-C, Nociceptin receptor,	
	K3 opiate receptor, Kappa-type 3 opioid receptor, Orphanin FQ receptor, Oprl1, Oor	
	Background: G-protein coupled opioid receptor that functions as receptor for the endogenous	
	neuropeptide nociceptin. Ligand binding causes a conformation change that triggers signaling	
	via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream	
	effectors. Signaling via G proteins mediates inhibition of adenylate cyclase activity and calcium	
	channel activity. Arrestins modulate signaling via G proteins and mediate the activation of	
	alternative signaling pathways that lead to the activation of MAP kinases. Plays a role in	
	modulating nociception and the perception of pain. Plays a role in the regulation of locomotor	
	activity by the neuropeptide nociceptin.	
Gene ID:	18389	
UniProt:	P35377	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	