

Datasheet for ABIN7260033

anti-PSMC2 antibody





Overview

Quantity:	200 μL
Target:	PSMC2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PSMC2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant fusion protein of human PSMC2 (NP_002794.1).
Isotype:	IgG
Characteristics:	Polyclonal Antibody
Purification:	Affinity purification

Target Details

Target:	PSMC2
Alternative Name:	PSMC2 (PSMC2 Products)
Background:	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure
	composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4
	rings of 28 non-identical subunits, 2 rings are composed of 7 alpha subunits and 2 rings are
	composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6

ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases which have a chaperone-like activity. This subunit has been shown to interact with several of the basal transcription factors so, in addition to participation in proteasome functions, this subunit may participate in the regulation of transcription. This subunit may also compete with PSMC3 for binding to the HIV tat protein to regulate the interaction between the viral protein and the transcription complex. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Molecular Weight:

Observed MW: 49 kDa

Calculated MW: 33 kDa/48 kDa

Gene ID:

5701

UniProt:

P35998

Pathways:

Mitotic G1-G1/S Phases, DNA Replication, Synthesis of DNA, Ubiquitin Proteasome Pathway

Application Details

Application Notes:

Storage Comment:

WB 1:500-1:2000 IHC 1:50-1:200 IF 1:50-1:200


Restrictions:

For Research Use only

Handling


Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS with 0.02 % sodium azide, 50 % glycerol, pH 7.3
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C

Store at -20°C. Avoid freeze / thaw cycles.




Immunofluorescence

Image 1. Immunofluorescence analysis of U2OS cells using PSMC2 Polyclonal Antibody



Western Blotting

Image 2. Western blot analysis of extracts of various cell lines using PSMC2 Polyclonal Antibody at dilution of 1:1000.



Immunohistochemistry (Paraffin-embedded Sections)

Image 3. Immunohistochemistry of paraffin-embedded Mouse heart using PSMC2 Polyclonal Antibody at dilution of 1:100 (40x lens).

Please check the product details page for more images. Overall 4 images are available for ABIN7260033.