
antibodies -online.com







anti-IGF2 antibody

2 Images



Go to Product page

Overview

Quantity:	200 μL
Target:	IGF2
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IGF2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant fusion protein of human IGF2 (NP_000603.1).
Isotype:	IgG
Characteristics:	Polyclonal Antibody
Purification:	Affinity purification

Target Details

Target:	IGF2
Alternative Name:	IGF2 (IGF2 Products)
Background:	This gene encodes a member of the insulin family of polypeptide growth factors, which are involved in development and growth. It is an imprinted gene, expressed only from the paternal allele, and epigenetic changes at this locus are associated with Wilms tumour, Beckwith-Wiedemann syndrome, rhabdomyosarcoma, and Silver-Russell syndrome. A read-through INS-

Target Details


	IGF2 gene exists, whose 5' region overlaps the INS gene and the 3' region overlaps this gene. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
Molecular Weight:	Observed_MW: 28 kDa Calculated_MW: 20 kDa/26 kDa
Gene ID:	3481
UniProt:	P01344
Pathways:	Hormone Activity, Regulation of Hormone Metabolic Process, Regulation of Hormone Biosynthetic Process, Regulation of Carbohydrate Metabolic Process, Activated T Cell Proliferation


Application Details

Application Notes:	WB 1:500-1:2000 IF 1:50-1:100
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS with 0.02 % sodium azide, 50 % glycerol, pH 7.3
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Avoid freeze / thaw cycles.





Immunofluorescence

Image 1. Immunofluorescence analysis of A-549 cells using IGF2 Polyclonal Antibody

Western Blotting

Image 2. Western blot analysis of extracts of various cell lines using IGF2 Polyclonal Antibody at dilution of 1:1000.