antibodies -online.com





anti-EXOSC4 antibody





Overview

Quantity:	200 μL
Target:	EXOSC4
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EXOSC4 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	Recombinant fusion protein of human EXOSC4 (NP_061910.1).
Isotype:	IgG
Characteristics:	Polyclonal Antibody
Purification:	Affinity purification

Target Details

l arget:	EXOSC4
Alternative Name:	EXOSC4 (EXOSC4 Products)
Background:	Non-catalytic component of the RNA exosome complex which has 3'->5' exoribonuclease
	activity and participates in a multitude of cellular RNA processing and degradation events. In
	the nucleus, the RNA exosome complex is involved in proper maturation of stable RNA species
	such as rRNA, snRNA and snoRNA, in the elimination of RNA processing by-products and non-

coding 'pervasive' transcripts, such as antisense RNA species and promoter-upstream transcripts (PROMPTs), and of mRNAs with processing defects, thereby limiting or excluding their export to the cytoplasm. The RNA exosome may be involved in Ig class switch recombination (CSR) and/or Ig variable region somatic hypermutation (SHM) by targeting AICDA deamination activity to transcribed dsDNA substrates. In the cytoplasm, the RNA exosome complex is involved in general mRNA turnover and specifically degrades inherently unstable mRNAs containing AU-rich elements (AREs) within their 3' untranslated regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs. It seems to be involved in degradation of histone mRNA. The catalytic inactive RNA exosome core complex of 9 subunits (Exo-9) is proposed to play a pivotal role in the binding and presentation of RNA for ribonucleolysis, and to serve as a scaffold for the association with catalytic subunits and accessory proteins or complexes. EXOSC4 binds to ARE-containing RNAs.

Molecular Weight:

Observed_MW: 26 kDa

Calculated_MW: 26 kDa

Gene ID:

54512

UniProt:

Q9NPD3


Application Details

Application Notes: WB 1:500-1:2000

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS with 0.02 % sodium azide, 50 % glycerol, pH 7.3
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Avoid freeze / thaw cycles.



Western Blotting

Image 1. Western blot analysis of extracts of various cell lines using EXOSC4 Polyclonal Antibody at dilution of 1:3000.