

Datasheet for ABIN7265216  
**anti-STAT3 antibody (pSer727)**[Go to Product page](#)

## 5 Images

## Overview

|                      |                                      |
|----------------------|--------------------------------------|
| Quantity:            | 200 µL                               |
| Target:              | STAT3                                |
| Binding Specificity: | pSer727                              |
| Reactivity:          | Human, Mouse, Rat                    |
| Host:                | Rabbit                               |
| Clonality:           | Polyclonal                           |
| Conjugate:           | This STAT3 antibody is un-conjugated |
| Application:         | Immunohistochemistry (IHC)           |

## Product Details

|                  |  |
|------------------|--|
| Immunogen:       | A synthetic phosphorylated peptide around S727 of human Stat3 (NP_644805.1). |
| Isotype:         | IgG  |
| Characteristics: | Phosphorylated antibody  |
| Purification:    | Affinity purification  |

## Target Details

|                   |  |
|-------------------|--|
| Target:           | STAT3  |
| Alternative Name: | Stat3 ( <a href="#">STAT3 Products</a> )   |
| Background:       | The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor |

## Target Details

associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. Mutations in this gene are associated with infantile-onset multisystem autoimmune disease and hyper-immunoglobulin E syndrome. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Gene ID: 6774

UniProt: [P40763](#)

Pathways: [JAK-STAT Signaling](#), [RTK Signaling](#), [Interferon-gamma Pathway](#), [Neurotrophin Signaling Pathway](#), [Dopaminergic Neurogenesis](#), [Response to Growth Hormone Stimulus](#), [Carbohydrate Homeostasis](#), [Stem Cell Maintenance](#), [Hepatitis C](#), [Protein targeting to Nucleus](#), [Feeding Behaviour](#), [CXCR4-mediated Signaling Events](#), [Signaling of Hepatocyte Growth Factor Receptor](#)

## Application Details

Application Notes: IHC 1:50-1:200

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 mg/mL


Buffer: PBS with 0.02 % sodium azide, 50 % glycerol, pH 7.3

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C

Storage Comment: Store at -20°C. Avoid freeze / thaw cycles.




#### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** Immunohistochemistry of paraffin-embedded Mouse lung using Phospho-Stat3(S727) Polyclonal Antibody at dilution of 1:100 (40x lens).



#### Immunohistochemistry (Paraffin-embedded Sections)

**Image 2.** Immunohistochemistry of paraffin-embedded Rat lung using Phospho-Stat3(S727) Polyclonal Antibody at dilution of 1:100 (40x lens).



#### Immunohistochemistry (Paraffin-embedded Sections)

**Image 3.** Immunohistochemistry of paraffin-embedded Human lung using Phospho-Stat3(S727) Polyclonal Antibody at dilution of 1:100 (40x lens).

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN7265216.