

Datasheet for ABIN7265937

**anti-BAZ2A antibody**

5 Images

[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	BAZ2A
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BAZ2A antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (IHC)

## Product Details

Purpose:	BAZ2A Rabbit pAb
Immunogen:	Recombinant fusion protein of Human BAZ2A.
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Characteristics:	Polyclonal Antibodies
Purification:	Affinity purification

## Target Details

Target:	BAZ2A
Alternative Name:	BAZ2A ( <a href="#">BAZ2A Products</a> )
Background:	Regulatory subunit of the ATP-dependent NoRC-1 and NoRC-5 ISWI chromatin remodeling

## Target Details

complexes, which form ordered nucleosome arrays on chromatin and facilitate access to DNA during DNA-templated processes such as DNA replication, transcription, and repair. Both complexes regulate the spacing of nucleosomes along the chromatin and have the ability to slide mononucleosomes to the center of a DNA template. Directly stimulates the ATPase activity of SMARCA5 in the NoRC-5 ISWI chromatin remodeling complex. The NoRC-1 ISWI chromatin remodeling complex has a lower ATP hydrolysis rate than the NoRC-5 ISWI chromatin remodeling complex. Within the NoRC-5 ISWI chromatin remodeling complex, mediates silencing of a fraction of rDNA by recruiting histone-modifying enzymes and DNA methyltransferases, leading to heterochromatin formation and transcriptional silencing (By similarity. In the complex, it plays a central role by being recruited to rDNA and by targeting chromatin modifying enzymes such as HDAC1, leading to repress RNA polymerase I transcription (By similarity. Recruited to rDNA via its interaction with TTF1 and its ability to recognize and bind histone H4 acetylated on 'Lys-16' (H4K16ac, leading to deacetylation of H4K5ac, H4K8ac, H4K12ac but not H4K16ac (By similarity. Specifically binds pRNAs, 150-250 nucleotide RNAs that are complementary in sequence to the rDNA promoter, pRNA-binding is required for heterochromatin formation and rDNA silencing (By similarity.,TIP5,WALp3,BAZ2A / TIP5,BAZ2A,Epigenetics & Nuclear Signaling,BAZ2A

Gene ID: 11176

UniProt: [Q9UIF9](#)

## Application Details

Application Notes: WB,1:500 - 1:2000,IHC,1:50 - 1:100,IF,1:50 - 1:200

Restrictions: For Research Use only

## Handling

Format: Liquid


Buffer: PBS with 0.02 % sodium azide,50 % glycerol, pH 7.3.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C

Storage Comment: Store at -20°C. Avoid freeze / thaw cycles.




### Immunofluorescence

**Image 1.** Immunofluorescence analysis of U-2 OS cells using BA antibody (ABIN7265937) at dilution of 1:100. Blue: DAPI for nuclear staining.



### Western Blotting

**Image 2.** Western blot analysis of extracts of various cell lines, using BA antibody (ABIN7265937) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 180s.



### Immunohistochemistry

**Image 3.** Immunohistochemistry of paraffin-embedded human mammary cancer using BA antibody (ABIN7265937) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN7265937.