



[Go to Product page](#)

Datasheet for ABIN7266399

anti-CFAP61 antibody

3 Images

Overview

Quantity:	100 µL
Target:	CFAP61
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CFAP61 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunofluorescence (IF)

Product Details

Purpose:	CFAP61 Rabbit pAb
Immunogen:	A synthetic peptide of human CFAP61.
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Characteristics:	Polyclonal Antibodies
Purification:	Affinity purification

Target Details

Target:	CFAP61
Alternative Name:	CFAP61 (CFAP61 Products)
Background:	May regulate cilium motility through its role in the assembly of the axonemal radial

Target Details

spokes.,C20orf26,CaM-IP3,dJ1002M8.3,dJ1178H5.4,CFAP61,Cell Biology & Developmental Biology,CFAP61

Gene ID: 26074

UniProt: [Q8NHU2](#)

Application Details

Application Notes: WB,1:500 - 1:2000,IHC,1:50 - 1:200,IF,1:50 - 1:200

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: PBS with 0.02 % sodium azide,50 % glycerol, pH 7.3.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C


Storage Comment: Store at -20°C. Avoid freeze / thaw cycles.

Images




Immunofluorescence

Image 1. Immunofluorescence analysis of Mouse testis using CF Rabbit pAb (ABIN7266399) at dilution of 1:100. Blue: DAPI for nuclear staining.



Western Blotting

Image 2. Western blot analysis of extracts of Mouse eye, using CF antibody (ABIN7266399) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 3 min.



Immunohistochemistry

Image 3. Immunohistochemistry of paraffin-embedded Human colon using CF Rabbit pAb (ABIN7266399) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.