



Datasheet for ABIN7267102
anti-FAM117A antibody



[Go to Product page](#)

5 Images

Overview

Quantity:	100 µL
Target:	FAM117A
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAM117A antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Purpose:	FAM117A Rabbit pAb
Immunogen:	Recombinant protein of human FAM117A.
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Characteristics:	Polyclonal Antibodies
Purification:	Affinity purification

Target Details

Target:	FAM117A
Alternative Name:	FAM117A (FAM117A Products)
Background:	FAM117A

Target Details

Molecular Weight: 48.3kDa

Gene ID: 81558

UniProt: [Q9C073](#)

Application Details

Application Notes: WB,1:1000 - 1:2000,IHC,1:50 - 1:200

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: PBS with 0.02 % sodium azide,50 % glycerol, pH 7.3.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C


Storage Comment: Store at -20°C. Avoid freeze / thaw cycles.

Images




Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded mouse liver using FA Rabbit pAb (ABIN7267102) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Western Blotting

Image 2. Western blot analysis of extracts of various cell lines, using FA antibody (ABIN7267102) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 3s.



Immunohistochemistry

Image 3. Immunohistochemistry of paraffin-embedded human thyroid cancer using FA Rabbit pAb (ABIN7267102) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN7267102.