



[Go to Product page](#)

Datasheet for ABIN7267116
anti-FAM3D antibody (AA 24-204)

5 Images

Overview

Quantity:	100 µL
Target:	FAM3D
Binding Specificity:	AA 24-204
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAM3D antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Purpose:	FAM3D Rabbit pAb
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 24-204 of human FAM3D (NP_620160.1).
Sequence:	RSYMSFSMKT IRLPRWLAAS PTKEIQVKKY KGLIKPCPA NYFAFKICSG AANVVGPTMC FEDRMIMSPV KNNVGRGLNI ALVNGTTGAV LGQKAFDMYS GDVMHLVKFL KEIPGGALVL VASYDDPGTK MNDESRKLFSDLGSSYAKQL GFRDSWVFIG AKDLRGKSPF EQFLKNSPDT N
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Characteristics:	Polyclonal Antibodies
Purification:	Affinity purification

Target Details

Target:	FAM3D
Alternative Name:	FAM3D (FAM3D Products)
Background:	EF7,OIT1,FAM3D,Immunology & Inflammation,Cytokines,FAM3D
Gene ID:	131177
UniProt:	Q96BQ1


Application Details

Application Notes:	WB,1:500 - 1:2000,IHC,1:50 - 1:200,IF,1:50 - 1:200
Restrictions:	For Research Use only

Handling



Format:	Liquid
Buffer:	PBS with 0.02 % sodium azide,50 % glycerol, pH 7.3.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Avoid freeze / thaw cycles.

Images



Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded Human colon using FD Rabbit pAb (ABIN7267116) at dilution of 1:50 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Western Blotting

Image 2. Western blot analysis of extracts of various cell lines, using FD antibody (ABIN7267116) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.

Immunohistochemistry

Image 3. Immunohistochemistry of paraffin-embedded Mouse testis using FD Rabbit pAb (ABIN7267116) at dilution of 1:50 (20x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN7267116.