



[Go to Product page](#)

Datasheet for ABIN7267216

## anti-FMO3 antibody

### 5 Images

#### Overview

Quantity:	100 µL
Target:	FMO3
Reactivity:	Human
Host:	Rabbit
Clonality:	Monoclonal
Conjugate:	This FMO3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF)

#### Product Details

Purpose:	FMO3 Rabbit mAb
Immunogen:	Recombinant protein of human FMO3.
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Characteristics:	Monoclonal Antibodies
Purification:	Affinity purification

#### Target Details

Target:	FMO3
Alternative Name:	FMO3 ( <a href="#">FMO3 Products</a> )
Background:	Flavin-containing monooxygenases (FMO) are an important class of drug-metabolizing

## Target Details

---

enzymes that catalyze the NADPH-dependent oxygenation of various nitrogen-,sulfur-, and phosphorous-containing xenobiotics such as therapeutic drugs, dietary compounds, pesticides, and other foreign compounds. The human FMO gene family is composed of 5 genes and multiple pseudogenes. FMO members have distinct developmental- and tissue-specific expression patterns. The expression of this FMO3 gene, the major FMO expressed in adult liver, can vary up to 20-fold between individuals. This inter-individual variation in FMO3 expression levels is likely to have significant effects on the rate at which xenobiotics are metabolised and, therefore, is of considerable interest to the pharmaceutical industry. This transmembrane protein localizes to the endoplasmic reticulum of many tissues. Alternative splicing of this gene results in multiple transcript variants encoding different isoforms. Mutations in this gene cause the disorder trimethylaminuria (TMAu) which is characterized by the accumulation and excretion of unmetabolized trimethylamine and a distinctive body odor. In healthy individuals, trimethylamine is primarily converted to the non odorous trimethylamine N-oxide.,FMO3, FMOII, TMAU, dJ127D3.1, flavin containing monooxygenase 3,Signal Transduction,Endocrine & Metabolism,Drug metabolism,FMO3

---

Molecular Weight: 56kDa

---

Gene ID: 2328

---

UniProt: [P31513](#)

## Application Details

---

Application Notes: WB,1:500 - 1:2000,IF,1:50 - 1:200

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Buffer: PBS with 0.02 % sodium azide,0.05 % BSA,50 % glycerol, pH 7.3.

---

Preservative: Sodium azide

---


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

---

Storage: -20 °C


---

Storage Comment: Store at -20°C. Avoid freeze / thaw cycles.




### Immunofluorescence

**Image 1.** Immunofluorescence analysis of Mouse liver cells using FMO3 antibody (ABIN7267216) at dilution of 1:100. Blue: DAPI for nuclear staining.



### Western Blotting

**Image 2.** Western blot analysis of extracts of various cell lines, using FMO3 antibody (ABIN7267216) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.



### Western Blotting

**Image 3.** Western blot analysis of extracts of various cell lines, using FMO3 antibody (ABIN7267216) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 5s.

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN7267216.