.-online.com antibodies

Datasheet for ABIN7267629 anti-HNRNPC antibody

2 Images



Overview

Quantity:	100 μL
Target:	HNRNPC
Reactivity:	Human
Host:	Rabbit
Clonality:	Monoclonal
Conjugate:	This HNRNPC antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF)

Product Details


Purpose:	hnRNP C Rabbit mAb
Immunogen:	A synthesized peptide derived from human hnRNP C
lsotype:	lgG
Cross-Reactivity:	Human, Mouse
Characteristics:	Monoclonal Antibodies
Purification:	Affinity purification


Target Details

Target:	HNRNPC
Alternative Name:	HNRNPC (HNRNPC Products)
Background:	This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7267629 | 04/24/2024 | Copyright antibodies-online. All rights reserved.

cleoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with geneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the as and appear to influence pre-mRNA processing and other aspects of mRNA metabolism ansport. While all of the hnRNPs are present in the nucleus, some seem to shuttle en the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding ties. The protein encoded by this gene can act as a tetramer and is involved in the ably of 40S hnRNP particles. Multiple transcript variants encoding at least two different ms have been described for this gene. [provided by RefSeq, Jul 2008],C1, C2, HNRNP, C, SNRPC,Epigenetics & Nuclear Signaling,RNA Binding,HNRNPC
as and appear to influence pre-mRNA processing and other aspects of mRNA metabolism ansport. While all of the hnRNPs are present in the nucleus, some seem to shuttle en the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding ties. The protein encoded by this gene can act as a tetramer and is involved in the ably of 40S hnRNP particles. Multiple transcript variants encoding at least two different ms have been described for this gene. [provided by RefSeq, Jul 2008],C1, C2, HNRNP, C, SNRPC,Epigenetics & Nuclear Signaling,RNA Binding,HNRNPC
ansport. While all of the hnRNPs are present in the nucleus, some seem to shuttle en the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding ties. The protein encoded by this gene can act as a tetramer and is involved in the ably of 40S hnRNP particles. Multiple transcript variants encoding at least two different ms have been described for this gene. [provided by RefSeq, Jul 2008],C1, C2, HNRNP, C, SNRPC,Epigenetics & Nuclear Signaling,RNA Binding,HNRNPC
en the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding ties. The protein encoded by this gene can act as a tetramer and is involved in the ably of 40S hnRNP particles. Multiple transcript variants encoding at least two different ms have been described for this gene. [provided by RefSeq, Jul 2008],C1, C2, HNRNP, C, SNRPC,Epigenetics & Nuclear Signaling,RNA Binding,HNRNPC
ties. The protein encoded by this gene can act as a tetramer and is involved in the ably of 40S hnRNP particles. Multiple transcript variants encoding at least two different ms have been described for this gene. [provided by RefSeq, Jul 2008],C1, C2, HNRNP, C, SNRPC,Epigenetics & Nuclear Signaling,RNA Binding,HNRNPC
bly of 40S hnRNP particles. Multiple transcript variants encoding at least two different ns have been described for this gene. [provided by RefSeq, Jul 2008],C1, C2, HNRNP, C, SNRPC,Epigenetics & Nuclear Signaling,RNA Binding,HNRNPC
ns have been described for this gene. [provided by RefSeq, Jul 2008],C1, C2, HNRNP, C, SNRPC,Epigenetics & Nuclear Signaling,RNA Binding,HNRNPC
C, SNRPC,Epigenetics & Nuclear Signaling,RNA Binding,HNRNPC
0
500 - 1:2000,IF,1:50 - 1:200
search Use only
ith 0.02 % sodium azide,0.05 % BSA,50 % glycerol, pH 7.3.
n azide
roduct contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
l be handled by trained staff only.





Immunofluorescence

Image 1. Immunofluorescence analysis of U-2 OS cells using hnRNP C Rabbit mAb (ABIN7267629) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

Western Blotting

Image 2. Western blot analysis of extracts of various cell lines, using hnRNP C Rabbit mAb (ABIN7267629) at 1:1000 dilution.Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution.Lysates/proteins: 25 µg per lane.Blocking buffer: 3 % nonfat dry milk in TBST.Detection: ECL Basic Kit (RM00020).Exposure time: 3s.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7267629 | 04/24/2024 | Copyright antibodies-online. All rights reserved.