



[Go to Product page](#)

Datasheet for ABIN7267736  
**anti-HIST1H2BK antibody**

4 Images

### Overview

Quantity:	100 µL
Target:	HIST1H2BK
Reactivity:	Human
Host:	Rabbit
Clonality:	Monoclonal
Conjugate:	This HIST1H2BK antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB)

### Product Details

Purpose:	Histone H2B Rabbit mAb
Immunogen:	A synthesized peptide derived from human Histone H2B
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Characteristics:	Monoclonal Antibodies
Purification:	Affinity purification

### Target Details

Target:	HIST1H2BK
Alternative Name:	HIST1H2BK ( <a href="#">HIST1H2BK Products</a> )
Background:	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the

## Target Details

---

chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-dependent histone that is a member of the histone H2B family. The protein encoded is an antimicrobial protein with antibacterial and antifungal activity. Two transcripts that encode the same protein have been identified for this gene, which is found in the histone microcluster on chromosome 6p21.33. [provided by RefSeq, Aug 2015],H2B/S, H2BFAiii, H2BFT, H2BK,Epigenetics & Nuclear Signaling,Histones,HIST1H2BK

---

Molecular Weight: 14kDa

---

Gene ID: 85236

---

UniProt: [O60814](#)

## Application Details

---

Application Notes: WB,1:500 - 1:2000,IHC,1:50 - 1:200

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Buffer: PBS with 0.02 % sodium azide,0.05 % BSA,50 % glycerol, pH 7.3.

---

Preservative: Sodium azide

---


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

---

Storage: -20 °C


---

Storage Comment: Store at -20°C. Avoid freeze / thaw cycles.




### Immunohistochemistry

**Image 1.** Immunohistochemistry of paraffin-embedded mouse kidney using Histone H2B Rabbit mAb (ABIN7267736) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



### Western Blotting

**Image 2.** Western blot analysis of extracts of various cell lines, using Histone H2B antibody (ABIN7267736) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 60s.



### Immunohistochemistry

**Image 3.** Immunohistochemistry of paraffin-embedded rat brain using Histone H2B Rabbit mAb (ABIN7267736) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN7267736.