



Datasheet for ABIN7267767  
**anti-HOXB4 antibody**



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	HOXB4
Reactivity:	Human
Host:	Rabbit
Clonality:	Monoclonal
Conjugate:	This HOXB4 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Purpose:	HOXB4 Rabbit mAb
Immunogen:	A synthesized peptide derived from human HOXB4
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Characteristics:	Monoclonal Antibodies
Purification:	Affinity purification

Target Details

Target:	HOXB4
Alternative Name:	HOXB4 ( <a href="#">HOXB4 Products</a> )
Background:	This gene is a member of the Antp homeobox family and encodes a nuclear protein with a

## Target Details

---

homeobox DNA-binding domain. It is included in a cluster of homeobox B genes located on chromosome 17. The encoded protein functions as a sequence-specific transcription factor that is involved in development. Intracellular or ectopic expression of this protein expands hematopoietic stem and progenitor cells in vivo and in vitro, making it a potential candidate for therapeutic stem cell expansion. [provided by RefSeq, Jul 2008],HOX-2.6, HOX2, HOX2F,Epigenetics & Nuclear Signaling,Hematopoietic Progenitors,Stem Cells,Transcription Factors,HOXB4

---

Molecular Weight: 34kDa

---

Gene ID: 3214

---

UniProt: [P17483](#)

## Application Details

---

---

Application Notes: WB,1:500 - 1:2000

---

Restrictions: For Research Use only

## Handling

---

---

Format: Liquid

---

Buffer: PBS with 0.02 % sodium azide,0.05 % BSA,50 % glycerol, pH 7.3.

---

Preservative: Sodium azide

---


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

---

Storage: -20 °C

---

Storage Comment: Store at -20°C. Avoid freeze / thaw cycles.



### Western Blotting

**Image 1.** Western blot analysis of extracts of various cell lines, using HOXB4 Rabbit mAb (ABIN7267767) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 180s.