# antibodies -online.com





## anti-NDRG1 antibody





#### Overview

Quantity:	100 μL
Target:	NDRG1
Reactivity:	Human
Host:	Rabbit
Clonality:	Monoclonal
Conjugate:	This NDRG1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

#### **Product Details**

Purpose:	NDRG1 Rabbit mAb
Immunogen:	A synthesized peptide derived from human NDRG1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Characteristics:	Monoclonal Antibodies
Purification:	Affinity purification

### Target Details

Target:	NDRG1
Alternative Name:	NDRG1 (NDRG1 Products)
Background:	This gene is a member of the N-myc downregulated gene family which belongs to the

alpha/beta hydrolase superfamily. The protein encoded by this gene is a cytoplasmic protein involved in stress responses, hormone responses, cell growth, and differentiation. The encoded protein is necessary for p53-mediated caspase activation and apoptosis. Mutations in this gene are a cause of Charcot-Marie-Tooth disease type 4D, and expression of this gene may be a prognostic indicator for several types of cancer. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, May 2012],CAP43, CMT4D, DRG-1, DRG1, GC4, HMSNL, NDR1, NMSL, PROXY1, RIT42, RTP, TARG1, TDD5,Angiogenesis,Cancer,Cardiovascular,Epigenetics & Nuclear Signaling,Invasion and Metastasis,Neurodegenerative Diseases,Neuroscience,Protein phosphorylation,Signal Transduction,NDRG1

Molecular Weight: 43kDa

Gene ID: 10397

UniProt: 092597


#### **Application Details**

Application Notes: WB,1:500 - 1:2000,IHC,1:50 - 1:200

Restrictions: For Research Use only


#### Handling

Format:	Liquid
Buffer:	PBS with 0.02 % sodium azide,0.05 % BSA,50 % glycerol, pH 7.3.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Avoid freeze / thaw cycles.




#### **Immunohistochemistry**

**Image 1.** Immunohistochemistry of paraffin-embedded rat kidney using NDRG1 Rabbit mAb (ABIN7268783) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



#### **Western Blotting**

Image 2. Western blot analysis of extracts of various cell lines, using NDRG1 Rabbit mAb (ABIN7268783) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 μg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.



#### **Immunohistochemistry**

**Image 3.** Immunohistochemistry of paraffin-embedded mouse kidney using NDRG1 Rabbit mAb (ABIN7268783) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

Please check the product details page for more images. Overall 4 images are available for ABIN7268783.