



Datasheet for ABIN7270361
anti-SNRNP40 antibody



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	SNRNP40
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SNRNP40 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Purpose:	SNRNP40 Rabbit pAb
Immunogen:	Recombinant protein of human SNRNP40.
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Characteristics:	Polyclonal Antibodies
Purification:	Affinity purification

Target Details

Target:	SNRNP40
Alternative Name:	SNRNP40 (SNRNP40 Products)
Background:	This gene encodes a component of the U5 small nuclear ribonucleoprotein (snRNP) particle.

Target Details

The U5 snRNP is part of the spliceosome, a multiprotein complex that catalyzes the removal of introns from pre-messenger RNAs. [provided by RefSeq, Jul 2008],40K, HPRP8BP, PRP8BP, PRPF8BP, SPF38, WDR57,SNRNP40

Gene ID: 9410

UniProt: [Q96D17](#)

Application Details

Application Notes: WB,1:500 - 1:2000

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: PBS with 0.02 % sodium azide,50 % glycerol, pH 7.3.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C


Storage Comment: Store at -20°C. Avoid freeze / thaw cycles.

Images



Western Blotting

Image 1. Western blot analysis of extracts of various cell lines, using SNRNP40 antibody (ABIN7270361) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.



Western Blotting

Image 2. Western blot analysis of extracts of various cell lines, using SNRNP40 antibody (ABIN7270361) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3 % nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90s.