
antibodies - online.com







anti-SUPT5H antibody





Overview

Quantity:	100 μL
Target:	SUPT5H
Reactivity:	Human
Host:	Rabbit
Clonality:	Monoclonal
Conjugate:	This SUPT5H antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF)

Product Details

Purpose:	SUPT5H/SPT5 Rabbit mAb
Immunogen:	Recombinant protein of human SUPT5H/SPT5.
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Characteristics:	Monoclonal Antibodies
Purification:	Affinity purification

Target Details

Target:	SUPT5H
Alternative Name:	SUPT5H (SUPT5H Products)
Background:	Component of the DRB sensitivity-inducing factor complex (DSIF complex, which regulates

mRNA processing and transcription elongation by RNA polymerase II. DSIF positively regulates mRNA capping by stimulating the mRNA quanylyltransferase activity of RNGTT/CAP1A. DSIF also acts cooperatively with the negative elongation factor complex (NELF complex to enhance transcriptional pausing at sites proximal to the promoter. Transcriptional pausing may facilitate the assembly of an elongation competent RNA polymerase II complex. DSIF and NELF promote pausing by inhibition of the transcription elongation factor TFIIS/S-II. TFIIS/S-II binds to RNA polymerase II at transcription pause sites and stimulates the weak intrinsic nuclease activity of the enzyme. Cleavage of blocked transcripts by RNA polymerase II promotes the resumption of transcription from the new 3' terminus and may allow repeated attempts at transcription through natural pause sites. DSIF can also positively regulate transcriptional elongation and is required for the efficient activation of transcriptional elongation by the HIV-1 nuclear transcriptional activator, Tat. DSIF acts to suppress transcriptional pausing in transcripts derived from the HIV-1 LTR and blocks premature release of HIV-1 transcripts at terminator sequences., SUPT5H, SPT5, SPT5H, Tat-CT1, transcription elongation factor SPT5, Epigenetics & Nuclear Signaling, Transcription Factors, RNA Binding, Cell Biology & Developmental Biology, Apoptosis, SUPT5H


Molecular Weight:	155kDa
Gene ID:	6829
UniProt:	O00267


Application Details


Application Notes:	WB,1:500 - 1:2000,IHC,1:50 - 1:200,IF,1:50 - 1:200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	PBS with 0.02 % sodium azide,0.05 % BSA,50 % glycerol, pH 7.3.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Avoid freeze / thaw cycles.







Immunohistochemistry

Image 1. Immunohistochemistry of paraffin-embedded human colon carcinoma using SUPT5H/SPT5 Rabbit mAb (ABIN7270878) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Western Blotting

Image 2. Western blot analysis of extracts of HeLa cells, using SUPT5H/SPT5 antibody (ABIN7270878) at 1:1000 dilution.Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution.Lysates/proteins: 25 µg per lane.Blocking buffer: 3 % nonfat dry milk in TBST.Detection: ECL Basic Kit (RM00020). Exposure time: 10s.

Immunofluorescence

Image 3. Immunofluorescence analysis of C6 cells using SUPT5H/SPT5 Rabbit mAb (ABIN7270878) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

Please check the product details page for more images. Overall 6 images are available for ABIN7270878.