
# antibodies - online.com







# anti-TNFRSF19 antibody





### Overview

Quantity:	100 μL
Target:	TNFRSF19
Reactivity:	Human
Host:	Rabbit
Clonality:	Monoclonal
Conjugate:	This TNFRSF19 antibody is un-conjugated
Application:	Western Blotting (WB)

# **Product Details**

Purpose:	TROY/TNFRSF19 Rabbit mAb
Immunogen:	Recombinant protein of human TROY/TNFRSF19.
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Characteristics:	Monoclonal Antibodies
Purification:	Affinity purification

# Target Details

Target:	TNFRSF19
Alternative Name:	TROY (TNFRSF19 Products)
Background:	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is

highly expressed during embryonic development. It has been shown to interact with TRAF family members, and to activate JNK signaling pathway when overexpressed in cells. This receptor is capable of inducing apoptosis by a caspase-independent mechanism, and it is thought to play an essential role in embryonic development. Alternatively spliced transcript variants encoding distinct isoforms have been described.,TNFRSF19, TAJ, TAJ-alpha, TRADE, TROY, TNF receptor superfamily member 19,Cancer,Signal Transduction,Cell Biology & Developmental Biology,Apoptosis,Growth factor,TNF,Immunology & Inflammation,NF-kB Signaling Pathway,Neuroscience,TROY

Molecular Weight: 46kDa

Gene ID: 55504

UniProt: Q9NS68


# **Application Details**

Application Notes: WB,1:500 - 1:2000

Restrictions: For Research Use only

# Handling

Format:	Liquid
Buffer:	PBS with 0.02 % sodium azide,0.05 % BSA,50 % glycerol, pH 7.3.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Avoid freeze / thaw cycles.



# **Western Blotting**

**Image 1.** Western blot analysis of extracts of various cell lines, using TROY/TNFRSF19 antibody (ABIN7271069) at 1:1000 dilution.Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution.Lysates/proteins: 25 μg per lane.Blocking buffer: 3 % nonfat dry milk in TBST.Detection: ECL Basic Kit (RM00020).Exposure time: 1s.