



Datasheet for ABIN7271144
anti-USP14 antibody



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	USP14
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This USP14 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Purpose:	USP14 Rabbit pAb
Immunogen:	A synthetic peptide of human USP14
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Characteristics:	Polyclonal Antibodies
Purification:	Affinity purification

Target Details

Target:	USP14
Alternative Name:	USP14 (USP14 Products)
Background:	This gene encodes a member of the ubiquitin-specific processing (UBP) family of proteases

Target Details

that is a deubiquitinating enzyme (DUB) with His and Cys domains. This protein is located in the cytoplasm and cleaves the ubiquitin moiety from ubiquitin-fused precursors and ubiquitylated proteins. Mice with a mutation that results in reduced expression of the ortholog of this protein are retarded for growth, develop severe tremors by 2 to 3 weeks of age followed by hindlimb paralysis and death by 6 to 10 weeks of age. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.,USP14,TGT,Epigenetics & Nuclear Signaling,Cell Biology & Developmental Biology,Ubiquitin,USP14

Molecular Weight: 52kDa/54kDa/56kDa

Gene ID: 9097

UniProt: [P54578](#)

Application Details

Application Notes: WB,1:500 - 1:2000

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: PBS with 0.02 % sodium azide,50 % glycerol, pH 7.3.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Avoid freeze / thaw cycles.



Western Blotting

Image 1. Western blot analysis of extracts of various cell lines, using USP14 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (ABIN1684268 and ABIN3020597) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 2 min.