antibodies -online.com







APOA2 Protein (AA 19-100) (Fc Tag)





Overview

Quantity:	100 μg
Target:	APOA2
Protein Characteristics:	AA 19-100
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This APOA2 protein is labelled with Fc Tag.

Product Details


Purpose:	Human APOA2 Protein
Sequence:	Ala19-Gln100
Characteristics:	Recombinant Human APOA2 Protein is expressed from HEK293 with hFc tag at the C-Terminus.It contains Ala19-Gln100.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 μm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.

Target Details

Target:	APOA2
Alternative Name:	APOA2 (APOA2 Products)


Target Details

3	
Background:	APOA2 is a protein implicated in triglyceride, fatty acid and glucose metabolism. In pigs, the APOA2 gene is located on pig chromosome 4 (SSC4) in a QTL region affecting fatty acid composition, fatness and growth traits.
Molecular Weight:	36.1 kDa. Due to glycosylation, the protein migrates to 40-50 kDa based on Tris-Bis PAGE result.
UniProt:	P02652
Pathways:	Regulation of Lipid Metabolism by PPARalpha, Production of Molecular Mediator of Immune Response, Negative Regulation of Transporter Activity, Lipid Metabolism
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months



SDS-PAGE

Image 1. Human APOA2 on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .



Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 2. The purity of Human APOA2 is greater than 95 % as determined by SEC-HPLC.