
antibodies -online.com







TNFSF9 Protein (AA 68-251) (Fc Tag)

3 Ima

Images



Go to Product page

()	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	$I \vee I$	ew

Quantity:	100 μg
Target:	TNFSF9
Protein Characteristics:	AA 68-251
Origin:	Cynomolgus
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TNFSF9 protein is labelled with Fc Tag.

Product Details


Purpose:	Cynomolgus 4-1BB Ligand/TNFSF9 Protein	
Sequence:	Arg68-Glu251	
Characteristics:	Recombinant Cynomolgus 4-1BB Ligand/TNFSF9 Protein is expressed from HEK293 with hFc tag at the C-Terminus.It contains Arg68-Glu251.	
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC	
Sterility:	0.22 μm filtered	
Endotoxin Level: Less than 1EU per µg by the LAL method.		
Biological Activity Comment:	Immobilized Cynomolgus/Rhesus macaque 4-1BB, hFc Tag at 0.5µg/ml (100µl/Well) on the	
	plate. Dose response curve for Cynomolgus 4-1BB Ligand, hFc Tag with the EC50 of 6.7ng/ml	
	determined by ELISA. See testing image for detail.	

Target Details


rarget Details		
Target:	TNFSF9	
Alternative Name:	4-1BB Ligand (TNFSF9 Products)	
Background:	The 4-1BBL is the high affinity ligand of 4-1BB, also known as CD137L or TNFSF9. 4-1BB ligand (4-1BBL) is an inducible molecule present on several APC types, including B cells, macrophages and DCs.4-1BB:4-1BBL pathway seems to amplify the existing costimulatory signals, even if the engagement of 4-1BB in the presence of a strong TCR signaling can induce IL-2 production in a CD28-independent manner.	
Molecular Weight:	46 kDa. Due to glycosylation, the protein migrates to 50-52 kDa based on Tris-Bis PAGE result.	
NCBI Accession:	XP_015296398	
Pathways:	Activated T Cell Proliferation, Cancer Immune Checkpoints	
Application Details		
Restrictions:	For Research Use only	
Handling		
Format:	Lyophilized	
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.	
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.	
Storage:	-20 °C,-80 °C	
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	
Expiry Date:	12 months	

Cynomolgus 4-1BB Ligand, hFc Tag ELISA

0.05µg Cynomolgus/Rhesus macaque 4-1BB, His Tag Per Well



Log Cynomolgus 4-1BB Ligand, hFc Tag Conc.(µg/ml)




ELISA

Image 1. Immobilized Cynomolgus/Rhesus macaque 4-1BB, hFc Tag at $0.5 \,\mu\text{g/mL}$ (100 $\mu\text{L/Well}$) on the plate. Dose response curve for Cynomolgus 4-1BB Ligand, hFc Tag with the EC50 of 6.7 ng/mL determined by ELISA.

Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 2. The purity of Cynomolgus 4-1BB Ligand is greater than 95 % as determined by SEC-HPLC.



SDS-PAGE

 $\label{lem:mage 3.} \textbf{End GE} \ \ \textbf{Mage 3.} \ \ \textbf{Cynomolgus 4-1BB Ligand on Tris-Bis PAGE under reduced condition}. \ \ \textbf{The purity is greater than 95\%}.$