antibodies -online.com







BCAM Protein (AA 32-548) (His tag)

Images



Overview

Quantity:	100 μg
Target:	BCAM
Protein Characteristics:	AA 32-548
Origin:	Cynomolgus
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This BCAM protein is labelled with His tag.

Product Details

Purpose:	Cynomolgus BCAM Protein
Sequence:	Glu32-Gly548
Characteristics:	Recombinant Cynomolgus BCAM Protein is expressed from HEK293 with His tag at the C-Terminus.It contains Glu32-Gly548.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 μm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.

Target Details


Target:	BCAM
Alternative Name:	BCAM (BCAM Products)

Target Details

Expiry Date:


12 months

rai got Botano	
Background:	Lutheran/basal cell adhesion molecule (Lu/BCAM) is a transmembrane adhesion molecule expressed by erythrocytes and endothelial cells that can interact with the extracellular matrix protein laminin-α5. In sickle cell disease, Lu/BCAM is thought to contribute to adhesion of sickle erythrocytes to the vascular wall, especially during vaso-occlusive crises.
Molecular Weight:	57.42 kDa. Due to glycosylation, the protein migrates to 70-80 kDa based on Tris-Bis PAGE result.
UniProt:	A0A7N9CA99
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
F : D :	40



SDS-PAGE

Image 1. Cynomolgus BCAM on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .



Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 2. The purity of Cynomolgus BCAM is greater than 95 % as determined by SEC-HPLC.