
antibodies - online.com







BCAM Protein (AA 26-541) (His tag)

Images



Overview

| Quantity: | 100 μg |
|-------------------------------|---|
| Target: | BCAM |
| Protein Characteristics: | AA 26-541 |
| Origin: | Mouse |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This BCAM protein is labelled with His tag. |

Product Details

| Purpose: | Mouse BCAM Protein |
|------------------|--|
| Sequence: | Glu26-Ala541 |
| Characteristics: | Recombinant Mouse BCAM Protein is expressed from HEK293 with His tag at the C-Terminus.It contains Glu26-Ala541. |
| Purity: | > 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC |
| Sterility: | 0.22 µm filtered |
| Endotoxin Level: | Less than 1EU per µg by the LAL method. |

Target Details


| Target: | BCAM |
|-------------------|----------------------|
| Alternative Name: | BCAM (BCAM Products) |

Target Details

Expiry Date:


12 months

| expressed by erythrocytes and endothelial cells that can interact with the extracellular in protein laminin-q5. In sickle cell disease, Lu/BCAM is thought to contribute to adhesion erythrocytes to the vascular wall, especially during vaso-occlusive crises. Molecular Weight: 57.95 kDa. Due to glycosylation, the protein migrates to 68-75 kDa based on Tris-Bis Presult. UniProt: Q9R069 Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 precommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22 pm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is accommended to the protein of th | rangerbetane | |
|--|---------------------|---|
| result. UniProt: Q9R069 Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µ recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is ac protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt.,-80 °C for 3-6 months after reconstitution.,2-8 °C for 2-7 days after reconstitution.,Recommend to aliquot the protein | Background: | Lutheran/basal cell adhesion molecule (Lu/BCAM) is a transmembrane adhesion molecule expressed by erythrocytes and endothelial cells that can interact with the extracellular matrix protein laminin-α5. In sickle cell disease, Lu/BCAM is thought to contribute to adhesion of sickle erythrocytes to the vascular wall, especially during vaso-occlusive crises. |
| Application Details Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µ recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is ac protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt.,-80 °C for 3-6 months after reconstitution.,2-8 °C for 2-7 days after reconstitution.,Recommend to aliquot the protein | Molecular Weight: | 57.95 kDa. Due to glycosylation, the protein migrates to 68-75 kDa based on Tris-Bis PAGE result. |
| Restrictions: For Research Use only Handling Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µ recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is ac protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt.,-80 °C for 3-6 months after reconstitution.,2-8 °C for 2-7 days after reconstitution.,Recommend to aliquot the protein | UniProt: | Q9R069 |
| Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µ recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is ac protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt.,-80 °C for 3-6 months after reconstitution.,2-8 °C for 2-7 days after reconstitution.,Recommend to aliquot the protein | Application Details | |
| Format: Lyophilized Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µ recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is ac protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein | Restrictions: | For Research Use only |
| Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µ recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is ac protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein | Handling | |
| recommended. Dissolve the lyophilized protein in distilled water. Buffer: Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is accommendate protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein | Format: | Lyophilized |
| protectant before lyophilization. Storage: -20 °C,-80 °C Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt.,-80 °C for 3-6 months after reconstitution.,2-8 °C for 2-7 days after reconstitution.,Recommend to aliquot the protein | Reconstitution: | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water. |
| Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein | Buffer: | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization. |
| reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein | Storage: | -20 °C,-80 °C |
| | Storage Comment: | -20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |



SDS-PAGE

Image 1. Mouse BCAM on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .



Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 2. The purity of Mouse BCAM is greater than $95\,\%$ as determined by SEC-HPLC.