# antibodies - online.com







## BST2 Protein (AA 49-163) (His tag)



#### Image



#### Overview

Quantity:	100 μg
Target:	BST2
Protein Characteristics:	AA 49-163
Origin:	Cynomolgus
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This BST2 protein is labelled with His tag.

#### **Product Details**


Purpose:	Cynomolgus BST2 Protein
Sequence:	Ile49-Ser163
Characteristics:	Recombinant Cynomolgus BST2 Protein is expressed from HEK293 with His tag at the N-Terminus.It contains Ile49-Ser163.
Purity:	> 95 % as determined by Tris-Bis PAGE
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.

#### **Target Details**

Target:	BST2
Alternative Name:	BST2 (BST2 Products)

### Target Details

rarget betails	
Background:	Interferon-induced BST2 (bone marrow stromal cell antigen 2) inhibits viral replication by tethering enveloped virions to the cell surface to restrict viral release and by inducing the NFKB-dependent antiviral immune response. BST2 expression was significantly increased during porcine epidemic diarrhea virus (PEDV) infection of Vero cells by IRF1 targeting its promoter. Both the BST2 and N protein interacted with the E3 ubiquitin ligase MARCHF8/MARCH8 and the cargo receptor.
Molecular Weight:	13.78 kDa. Due to glycosylation, the protein migrates to 15-30 kDa based on Tris-Bis PAGE result.
NCBI Accession:	XP_005588438
Pathways:	Regulation of Leukocyte Mediated Immunity, Production of Molecular Mediator of Immune Response
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months



#### **SDS-PAGE**

**Image 1.** Cynomolgus BST2 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.