

Datasheet for ABIN7273992

**BST2 Protein (AA 49-161) (His tag)****3** Images[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	BST2
Protein Characteristics:	AA 49-161
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This BST2 protein is labelled with His tag.

## Product Details

Purpose:	Human BST2 Protein
Sequence:	Asn49-Ser161
Characteristics:	Recombinant Human BST2 Protein is expressed from HEK293 with His tag at the N-Terminus. It contains Asn49-Ser161.
Purity:	> 95 % as determined by Tris-Bis PAGE
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.
Biological Activity Comment:	Immobilized Human BST2, His Tag at 2µg/ml (100µl/Well) on the plate. Dose response curve for Anti-BST2 Antibody, hFc Tag with the EC50 of 15.9ng/ml determined by ELISA. See testing image for detail.

## Target Details


Target:	BST2
Alternative Name:	BST2 ( <a href="#">BST2 Products</a> )
Background:	Interferon-induced BST2 (bone marrow stromal cell antigen 2) inhibits viral replication by tethering enveloped virions to the cell surface to restrict viral release and by inducing the NFkB-dependent antiviral immune response. BST2 expression was significantly increased during porcine epidemic diarrhea virus (PEDV) infection of Vero cells by IRF1 targeting its promoter. Both the BST2 and N protein interacted with the E3 ubiquitin ligase MARCHF8/MARCH8 and the cargo receptor.
Molecular Weight:	14.5 kDa. Due to glycosylation, the protein migrates to 25-35 kDa based on Tris-Bis PAGE result.
Pathways:	<a href="#">Regulation of Leukocyte Mediated Immunity</a> , <a href="#">Production of Molecular Mediator of Immune Response</a>

## Application Details

Restrictions:	For Research Use only
---------------	-----------------------


## Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS ( pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months



SDS-PAGE


**Image 1.** Human BST2 on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .



Size-exclusion chromatography-High Pressure Liquid Chromatography

**Image 2.** The purity of Human BST2 is greater than 95 % as determined by SEC-HPLC.

**Human BST2, His Tag ELISA**  
0.2µg Human BST2, His Tag Per Well



ELISA

**Image 3.** Immobilized Human BST2, His Tag at 2 µg/mL (100 µL/Well) on the plate. Dose response curve for Anti-BST2 Antibody, hFc Tag with the EC50 of 15.9 ng/mL determined by ELISA.