

Datasheet for ABIN7274001

BTN3A1 Protein (AA 1-219) (His-Avi Tag)**3** Images[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	BTN3A1
Protein Characteristics:	AA 1-219
Origin:	Cynomolgus
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This BTN3A1 protein is labelled with His-Avi Tag.

Product Details

Purpose:	Cynomolgus BTN3A1/CD277 Protein
Sequence:	Gln1-Trp219
Characteristics:	Recombinant Cynomolgus BTN3A1/CD277 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Gln1-Trp219.
Purity:	> 95 % as determined by Tris-Bis PAGE, > 95 % as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.
Biological Activity Comment:	Immobilized Cynomolgus BTN3A1, His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-BTN3A1 Antibody, hFc Tag with the EC50 of 10.9ng/ml determined by ELISA. See testing image for detail.

Target Details

Target:	BTN3A1
Alternative Name:	BTN3A1 (BTN3A1 Products)
Background:	The three butyrophilin BTN3A molecules, BTN3A1, BTN3A2, and BTN3A3, are members of the B7/butyrophilin-like group of Ig superfamily receptors, which modulate the function of T cells. BTN3A1 controls activation of human V γ 9/V δ 2 T cells by direct or indirect presentation of self and nonself phosphoantigens (pAg).
Molecular Weight:	26.5 kDa. Due to glycosylation, the protein migrates to 30-35 kDa based on Tris-Bis PAGE result.
UniProt:	A0A330KWZ1
Pathways:	Activated T Cell Proliferation

Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22 μ m filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 $^{\circ}$ C,-80 $^{\circ}$ C
Storage Comment:	-20 to -80 $^{\circ}$ C for 12 months as supplied from date of receipt.,-80 $^{\circ}$ C for 3-6 months after reconstitution.,2-8 $^{\circ}$ C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months

Size-exclusion chromatography-High Pressure Liquid Chromatography



Image 1. The purity of Cynomolgus BTN3A1 is greater than 95 % as determined by SEC-HPLC.

SDS-PAGE

Image 2. Cynomolgus BTN3A1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .

ELISA

Image 3. Immobilized Cynomolgus BTN3A1, His Tag at 0.5 µg/mL (100 µL/Well) on the plate. Dose response curve for Anti-BTN3A1 Antibody, hFc Tag with the EC50 of 10.9 ng/mL determined by ELISA.



Cynomolgus BTN3A1, His Tag ELISA
0.05µg Cynomolgus BTN3A1, His Tag Per Well

