antibodies -online.com







IL17RA Protein (AA 33-320) (Fc Tag)





Overview

Quantity:	100 μg
Target:	IL17RA
Protein Characteristics:	AA 33-320
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL17RA protein is labelled with Fc Tag.

Product Details

Purpose:	Human IL-17R alpha/CD217 Protein
Sequence:	Leu33-Trp320
Characteristics:	Recombinant Human IL-17R alpha/CD217 Protein is expressed from HEK293 with hFc tag at the C-Terminus.It contains Leu33-Trp320.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.
Biological Activity Comment:	Immobilized Human IL-17A, His Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve
	for Human IL-17R alpha, hFc Tag with the EC50 of 78.0ng/ml determined by ELISA. The affinity
	constant of 2.18 nM as determined in SPR assay (Biacore T200). See testing image for detail.

Target Details


Target:	IL17RA
Alternative Name:	IL-17R alpha (IL17RA Products)
Background:	Interleukin 17 (also known as CTLA-8) is a T cell-expressed pleotropic cytokine. IL-17 binds to IL-17 receptor (IL-17 R) which shares no homology with any known family of receptors. While the expression of IL-17 is restricted to activated T cells, the IL-17 R mRNA exhibits a broad tissue distribution, and has been detected in virtually all cells and tissues tested. The human IL-17 R gene was localized to chromosome 22.
Molecular Weight:	60.2 kDa. Due to glycosylation, the protein migrates to 75-90 kDa based on Tris-Bis PAGE result.
Pathways:	SARS-CoV-2 Protein Interactome

Application Details

Restrictions: For Research Use only


Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months




Surface Plasmon Resonance

Image 1. Human IL-17R alpha, hFc Tag captured on CM5 Chip via protein A can bind Biotinylated Human IL-17A, His-Avi Tag with an affinity constant of 2.18 nM as determined in SPR assay (Biacore T200).



Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 2. The purity of Human IL-17R alpha/CD217 is greater than 95 % as determined by SEC-HPLC.



SDS-PAGE

 $\label{lemage 3. Human IL-17R alpha/CD217 on Tris-Bis PAGE} \\$ under reduced condition. The purity is greater than 95 % .

Please check the product details page for more images. Overall 4 images are available for ABIN7274230.