

Datasheet for ABIN7274287

**FCGR2B Protein (AA 46-224) (His tag)****3** Images[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	FCGR2B
Protein Characteristics:	AA 46-224
Origin:	Cynomolgus
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This FCGR2B protein is labelled with His tag.

## Product Details

Purpose:	Cynomolgus Fc gamma RIIB Protein
Sequence:	Ala46-Pro224
Characteristics:	Recombinant Cynomolgus Fc gamma RIIB Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Ala46-Pro224.
Purity:	> 95 % as determined by Tris-Bis PAGE, > 95 % as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.
Biological Activity Comment:	The affinity constant of 2.72 µM as determined in SPR assay (Biacore T200). See testing image for detail.

## Target Details

Target:	FCGR2B
Alternative Name:	Fc gamma RIIB ( <a href="#">FCGR2B Products</a> )
Background:	The Fc gamma Rs have been divided into three classes based on close relationships in their extracellular domains, these groups are designated Fc gamma RI (also known as CD64), Fc gamma RII (CD32), and Fc gamma RIII (CD16). Each group may be encoded by multiple genes and exist in different isoforms depending on species and cell type. The CD64 proteins are high affinity receptors ( $\sim 10^{-8}$ - $10^{-9}$ M) capable of binding monomeric IgG, whereas the CD16 and CD32 proteins bind IgG with lower affinities ( $\sim 10^{-6}$ - $10^{-7}$ M) only recognizing IgG aggregates surrounding multivalent antigens.
Molecular Weight:	21 kDa. Due to glycosylation, the protein migrates to 33-40 kDa based on Tris-Bis PAGE result.
UniProt:	<a href="#">Q8SPW3</a>
Pathways:	<a href="#">Cellular Response to Molecule of Bacterial Origin</a> , <a href="#">Regulation of Leukocyte Mediated Immunity</a> , <a href="#">Production of Molecular Mediator of Immune Response</a> , <a href="#">BCR Signaling</a>

## Application Details


Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS ( pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months


Surface Plasmon Resonance

**Image 1.** Trastuzumab captured on CM5 Chip via Protein A can bind Cynomolgus Fc gamma RIIB, His Tag with an affinity constant of 2.72  $\mu$ M as determined in SPR assay (Biacore T200).



Size-exclusion chromatography-High Pressure Liquid Chromatography

**Image 2.** The purity of Cynomolgus Fc gamma RIIB is greater than 95 % as determined by SEC-HPLC.



SDS-PAGE

**Image 3.** Cynomolgus Fc gamma RIIB on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .

