

Datasheet for ABIN7274416

CRTAM Protein (AA 18-287) (His tag)**2** Images[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	CRTAM
Protein Characteristics:	AA 18-287
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CRTAM protein is labelled with His tag.

Product Details

Purpose:	Human CRTAM Protein
Sequence:	Ser18-Gly287
Characteristics:	Recombinant Human CRTAM Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Ser18-Gly287.
Purity:	> 95 % as determined by Tris-Bis PAGE
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.
Biological Activity Comment:	Immobilized Human CRTAM, His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-CRTAM Antibody, hFc Tag with the EC50 of 57.3ng/ml determined by ELISA. See testing image for detail.

Target Details

Target:	CRTAM
Alternative Name:	CRTAM (CRTAM Products)
Background:	Class-I Restricted T Cell-Associated Molecule (CRTAM) is a protein that is expressed after T cell activation. The interaction of CRTAM with its ligand, nectin-like 2 (Nect2), is required for the efficient production of IL-17, IL-22, and IFN γ by murine CD4 T cells, and it plays a role in optimal CD8 T and NK cell cytotoxicity. CRTAM promotes the pro-inflammatory cytokine profile, therefore, it may take part in the immunopathology of autoimmune diseases such as diabetes type 1 or colitis.
Molecular Weight:	31.1 kDa. Due to glycosylation, the protein migrates to 55-70 kDa based on Tris-Bis PAGE result.
Pathways:	Regulation of Leukocyte Mediated Immunity , Positive Regulation of Immune Effector Process , Activated T Cell Proliferation

Application Details


Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22 μ m filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months


Human CRTAM, His Tag ELISA

0.05µg Human CRTAM, His Tag Per Well



ELISA

Image 1. Immobilized Human CRTAM, His Tag at 0.5 µg/mL (100 µL/Well) on the plate. Dose response curve for Anti-CRTAM Antibody, hFc Tag with the EC50 of 57.3 ng/mL determined by ELISA.



SDS-PAGE

Image 2. Human CRTAM on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .