antibodies -online.com







CXCL16 Protein (AA 30-205) (Fc Tag)



Image



Overview

| Quantity: | 100 μg |
|-------------------------------|--|
| Target: | CXCL16 |
| Protein Characteristics: | AA 30-205 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This CXCL16 protein is labelled with Fc Tag. |

Product Details

| Purpose: | Human CXCL16 Protein |
|------------------|--|
| Sequence: | Asn30-Thr205 |
| Characteristics: | Recombinant Human CXCL16 Protein is expressed from HEK293 with hFc tag at the C-Terminus.It contains Asn30-Thr205. |
| Purity: | > 95 % as determined by Tris-Bis PAGE |
| Sterility: | 0.22 µm filtered |
| Endotoxin Level: | Less than 1EU per µg by the LAL method. |

Target Details


| Target: | CXCL16 |
|-------------------|--------------------------|
| Alternative Name: | CXCL16 (CXCL16 Products) |

Target Details

Expiry Date:

12 months

| rai get Betaile | |
|---------------------|---|
| Background: | CXC chemokine ligand 16 (CXCL16) is a CXC soluble chemokine, an adhesion molecule and a cell surface scavenger receptor. CXCL16 regulates inflammation, tissue injury and fibrosis. Parenchymal renal cells, vascular wall cells, leukocytes and platelets express and/or release CXCL16 under the regulation of inflammatory mediators. CXCL16 expression is increased in experimental and human nephropathies. Targeting CXCL16 protected from experimental glomerular injury or interstitial fibrosis. |
| Molecular Weight: | 45.9 kDa. Due to glycosylation, the protein migrates to 70-75 kDa based on Tris-Bis PAGE result. |
| UniProt: | Q9H2A7 |
| Application Details | |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Lyophilized |
| Reconstitution: | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water. |
| Buffer: | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | -20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |
| F . D . | |



SDS-PAGE

Image 1. Human CXCL16 on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .