# antibodies - online.com







# **DPP4 Protein (His tag)**





#### Overview

| Quantity:                     | 100 μg                                      |
|-------------------------------|---|
| Target:                       | DPP4  |
| Origin:                       | Human                                       |
| Source:                       | HEK-293 Cells                               |
| Protein Type:                 | Recombinant                                 |
| Purification tag / Conjugate: | This DPP4 protein is labelled with His tag. |

## **Product Details**

| Sequence:                    | Asn29-Pro766   |
|------------------------------|--|
| Purity:                      | > 95% as determined by Tris-Bis PAGE,> 95% as determined by HPLC                             |
| Sterility:                   | 0.22 μm filtered   |
| Endotoxin Level:             | Less than 1EU per μg by the LAL method.  |
| Biological Activity Comment: | Immobilized Human DPPIV at 1µg/ml (100µl/well) on the plate. Dose response curve for Anti-   |
|                              | DPPIV Antibody, hFc Tag with the EC50 of 8.1ng/ml determined by ELISA. See testing image for |
|                              | detail.  |

# Target Details


| Target:           | DPP4  |
|-------------------|---|
| Alternative Name: | DPPIV (DPP4 Products)   |
| Background:       | ADCP2, CD26, DPPIV, DPP4, ADABP, TP103, EC 3.4.14.5,CD26/dipeptidyl peptidase (DPP)IV is a    |
|                   | membrane-bound protein found in many cell types of the body, and a soluble form is present in |

# **Target Details**

|                     | body fluids. There is longstanding evidence that various primary tumors and also metastases express DPPIV/CD26 to a variable extent.  |
|---------------------|---|
| Molecular Weight:   | 90.5 kDa. Due to glycosylation, the protein migrates to 91-115 kDa based on Tris-Bis PAGE result.   |
| UniProt:            | P27487  |
| Pathways:           | Peptide Hormone Metabolism, Regulation of Leukocyte Mediated Immunity   |
| Application Details |   |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Lyophilized   |
| Reconstitution:     | Centrifuge tubes before opening. Reconstituting to a concentration more than 100 $\mu$ g/mL is recommended (usually we use 1 mg/mL solution for lyophilization). Dissolve the lyophilized protein in distilled water. |
| Buffer:             | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 5 % trehalose is added as protectant before lyophilization.   |
| Storage:            | 4 °C,-80 °C   |
| Storage Comment:    | Reconstituted protein stable at -80°C for 12 months, 4°C for 1 week. Use a manual defrost freezer and avoid repeated freeze-thaw cycles.  |
| Expiry Date:        | 12 months   |

#### Human DPPIV, His ELISA

0.1µg Human DPPIV, His Tag Per Well




#### **ELISA**

**Image 1.** Immobilized Human DPPIV at  $1 \mu g/mL$  ( $100 \mu$  L/well) on the plate. Dose response curve for Anti-DPPIV Antibody, hFc Tag with the EC50 of 8.1 ng/mL determined by ELISA.

### **SDS-PAGE**

**Image 2.** Human DPPIV on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



# Size-exclusion chromatography-High Pressure Liquid Chromatography

**Image 3.** The purity of Human DPPIV Protein is greater than 95 % as determined by SEC-HPLC.

