

Datasheet for ABIN7274645

**Fc gamma RII (CD32) (AA 17-208) protein (His tag)****3** Images[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	Fc gamma RII (CD32)
Protein Characteristics:	AA 17-208
Origin:	Cynomolgus
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	His tag

## Product Details

Purpose:	Cynomolgus Fc gamma RIII/CD16 Protein
Sequence:	Gly17-Gln208
Characteristics:	Recombinant Cynomolgus Fc gamma RIII/CD16 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Gly17-Gln208.
Purity:	> 95 % as determined by Tris-Bis PAGE, > 95 % as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.
Biological Activity Comment:	The affinity constant of 0.251 µM as determined in SPR assay (Biacore T200). See testing image for detail.

## Target Details


Target:	Fc gamma RII (CD32)
Alternative Name:	Fc gamma RIII ( <a href="#">CD32 Products</a> )
Background:	Immunoglobulin G (IgG) Fc receptors play a critical role in linking IgG antibody-mediated immune responses with cellular effector functions. A high resolution map of the binding site on human IgG1 for human Fc gamma RI, Fc gamma RIIA, Fc gamma RIIB, Fc gamma RIIIA, and FcRn receptors has been determined. A common set of IgG1 residues is involved in binding to all Fc gamma R, Fc gamma RII and Fc gamma RIII also utilize residues outside this common set.
Molecular Weight:	23.1 kDa. Due to glycosylation, the protein migrates to 45-55 kDa based on Tris-Bis PAGE result.

## Application Details

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22µm filtered solution in PBS ( pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months



SDS-PAGE

**Image 1.** Cynomolgus Fc gamma RIII on Tris-Bis PAGE under reduced condition. The purity is greater than 95 % .

Size-exclusion chromatography-High Pressure Liquid Chromatography

**Image 2.** The purity of Cynomolgus Fc gamma RIII is greater than 95 % as determined by SEC-HPLC.

Surface Plasmon Resonance

**Image 3.** Rituximab captured on CM5 Chip via Protein A can bind Cynomolgus Fc gamma RIII, His Tag with an affinity constant of 0.251  $\mu$ M as determined in SPR assay (Biacore T200).